

FORM PTO-1082

Date: August 10, 1999
Docket No.: 06171.105005

ASSISTANT COMMISSIONER FOR PATENTS
Washington, D.C. 20231

Sir:

Transmitted herewith for filing is the patent application of:

Inventors: Gilles Gosselin, Jean-Louis Imbach, and Martin L. Bryant

For: β -L-2'-Deoxy-Nucleosides for the Treatment of Hepatitis B Virus

Enclosed are:

☒ Patent Application consisting of 51 pages of specification, 5 pages of claims, 1 page of abstract, and 3 sheets of drawings;

☐ Assignment Recordation Cover Sheet

☐ Assignment of the invention to _____

☐ Check No. _____ in the amount of \$40.00 in payment of Assignment Recordation

☐ A Declaration (unexecuted)

☒ A Declaration/Power of Attorney

☐ A verified statement to establish small entity status under 37 CFR 1.9 and 37 CFR 1.27

☒ Express Mail Transmittal Letter for Patent Application and Certificate of Mailing (Express Mail Label No. EH010184723US).

The filing fee has been calculated as shown below:

	Col. 1	Col. 2	Small/Large Ent.	
For:	No. Filed	No. Extra	Rate	Filing Fee
Basic Fee	Excess of Twenty Independent Claims (Excess of Three)		\$11.00/\$22.00 \$41.00/\$82.00	\$380.00/\$760.00
Total Claims	<u>12</u> - 20		x 18/9 =	
Independent Claims	<u>4</u> - 3	<u>1</u>	x 78/39 =	<u>78.00</u>
Multiple Dependent Claims			x 260/130 =	
			Total	<u>838.00</u>

* If the difference in Col. 1 is less than zero, enter "0" in Col. 2.

X Please charge Deposit Account No. 11-0980 in the amount of \$ 838.00. A duplicate of this sheet is enclosed.

 A check in the amount of \$ is enclosed. (Check No(s).)

X The Commissioner is hereby authorized to charge payment of the following fees associated with this communication or credit any overpayment to Deposit Account No. 11-0980. A duplicate copy of this sheet is enclosed.

X Any additional filing fees required under 37 CFR 1.16.

X Any patent application processing fees under 37 CFR 1.17.

Respectfully submitted,

By: Jacqueline Haley
Jacqueline Haley
Reg. No.: 41,457

CERTIFICATE OF MAILING

I HEREBY CERTIFY THAT THIS DOCUMENT IS BEING FORWARDED TO THE ASSISTANT COMMISSIONER FOR PATENTS, WASHINGTON, D.C. 20231, VIA UNITED STATES POSTAL SERVICE AS "EXPRESS MAIL" POST OFFICE TO ADDRESSEE SERVICE UNDER 37 CFR 1.10, LABEL NO. EH010184723US ON THIS 10TH DAY OF August, 1999.

Jacqueline Haley
Jacqueline Haley

β -L-2'-Deoxy-Nucleosides for the Treatment of Hepatitis B

This application claims priority to U.S. provisional application U.S.S.N. 60/096,110, filed on August 10, 1998 and U.S.S.N. 60/131,352, filed on April 28, 1999.

5

Background of the Invention

This invention is in the area of methods for the treatment of hepatitis B virus (also referred to as "HBV") that includes administering to a host in need thereof, either alone or in
10 combination, an effective amount of one or more of the active compounds disclosed herein, or a pharmaceutically acceptable prodrug or salt of one of these compounds.

HBV is second only to tobacco as a cause of human cancer. The mechanism by which HBV induces cancer is unknown, although it is postulated that it may directly trigger tumor
15 development, or indirectly trigger tumor development through chronic inflammation, cirrhosis, and cell regeneration associated with the infection.

Hepatitis B virus has reached epidemic levels worldwide. After a two to six month incubation period in which the host is unaware of the infection, HBV infection can lead to acute hepatitis and liver damage, that causes abdominal pain, jaundice, and elevated blood levels of
20 certain enzymes. HBV can cause fulminant hepatitis, a rapidly progressive, often fatal form of the disease in which massive sections of the liver are destroyed.

Patients typically recover from acute hepatitis. In some patients, however, high levels of viral antigen persist in the blood for an extended, or indefinite, period, causing a chronic infection. Chronic infections can lead to chronic persistent hepatitis. Patients infected with
25 chronic persistent HBV are most common in developing countries. By mid-1991, there were approximately 225 million chronic carriers of HBV in Asia alone, and worldwide, almost 300 million carriers. Chronic persistent hepatitis can cause fatigue, cirrhosis of the liver, and hepatocellular carcinoma, a primary liver cancer.

In western industrialized countries, high risk groups for HBV infection include those in contact with HBV carriers or their blood samples. The epidemiology of HBV is very similar to
30 that of acquired immune deficiency syndrome (AIDS), which accounts for why HBV infection is

common among patients with AIDS or AIDS related complex. However, HBV is more contagious than HIV.

However, more recently, vaccines have also been produced through genetic engineering and are currently used widely. Unfortunately, vaccines cannot help those already infected with HBV. Daily treatments with α -interferon, a genetically engineered protein, has also shown promise, but this therapy is only successful in about one third of treated patients. Further, interferon cannot be given orally.

A number of synthetic nucleosides have been identified which exhibit activity against HBV. The (-)-enantiomer of BCH-189, known as 3TC, claimed in U. S. Patent 5,539,116 to Liotta, et al., has been approved by the U.S. Food and Drug Administration for the treatment of hepatitis B. See also EPA 0 494 119 A1 filed by BioChem Pharma, Inc.

Cis-2-hydroxymethyl-5-(5-fluorocytosin-1-yl)-1,3-oxathiolane ("FTC") exhibits activity against HBV. See WO 92/15308; Furman, et al., "The Anti-Hepatitis B Virus Activities, Cytotoxicities, and Anabolic Profiles of the (-) and (+) Enantiomers of cis-5-Fluoro-1-[2-(Hydroxymethyl)-1,3-oxathiolane-5-yl]-Cytosine" Antimicrobial Agents and Chemotherapy, December 1992, page 2686-2692; and Cheng, et al., Journal of Biological Chemistry, Volume 267(20), 13938-13942 (1992).

von Janta-Lipinski et al. disclose the use of the L-enantiomers of 3'-fluoro-modified β -2'-deoxyribonucleoside 5'-triphosphates for the inhibition of hepatitis B polymerases (J. Med. Chem., 1998, 41,2040-2046). Specifically, the 5'-triphosphates of 3'-deoxy-3'-fluoro- β -L-thymidine (β -L-FTTP), 2',3'-dideoxy-3'-fluoro- β -L-cytidine (β -L-FdCTP), and 2',3'-dideoxy-3'-fluoro- β -L-5-methylcytidine (β -L-FMethCTP) were disclosed as effective inhibitors of HBV DNA polymerases.

WO 96/13512 to Genencor International, Inc. and Lipitek, Inc. discloses that certain L-ribofuranosyl nucleosides can be useful for the treatment of cancer and viruses. Specifically disclosed is the use of this class of compounds for the treatment of cancer and HIV.

U. S. Patent Nos. 5,565,438, 5,567,688 and 5,587,362 (Chu, et al.) disclose the use of 2'-fluoro-5-methyl- β -L-arabinofuranolyluridine (L-FMAU) for the treatment of hepatitis B and Epstein Barr virus.

Yale University and University of Georgia Research Foundation, Inc. disclose the use of L-FddC (β -L-5-fluoro-2',3'-dideoxycytidine) for the treatment of hepatitis B virus in WO 92/18517.

The synthetic nucleosides β -L-2'-deoxycytidine (β -L-2'-dC), β -L-2'-deoxythymidine (β -L-dT) and β -L-2'-deoxyadenosine (β -L-2'-dA), are known in the art. Antonin Holy first disclosed β -L-dC and β -L-dT in 1972, "Nucleic Acid Components and Their Analogs. CLIII. Preparation of 2'-deoxy-L-Ribonucleosides of the Pyrimidine Series," *Collect. Czech. Chem. Commun.* (1972), 37(12), 4072-87. Morris S. Zedeck et al. first disclosed β -L-dA for the inhibition of the synthesis of induced enzymes in *Pseudomonas testosteroni*, *Mol. Phys.* (1967), 3(4), 386-95.

Certain 2'-deoxy- β -L-erythro-pentofuranonucleosides are known to have antineoplastic and selected antiviral activities. Verri et al. disclose the use of 2'-deoxy- β -L-erythro-pentofuranonucleosides as antineoplastic agents and as anti-herpetic agents (*Mol. Pharmacol.* (1997), 51(1), 132-138 and *Biochem. J.* (1997), 328(1), 317-20). Saneyoshi et al. demonstrate the use of 2'-deoxy-L-ribonucleosides as reverse transcriptase (I) inhibitors for the control of retroviruses and for the treatment of AIDS, Jpn. Kokai Tokkyo Koho JP06293645 (1994).

Giovanni et al. tested 2'-deoxy- β -L-erythro-pentofuranonucleosides against partially pseudorabies virus (PRV), *Biochem. J.* (1993), 294(2), 381-5.

Chemotherapeutic uses of 2'-deoxy- β -L-erythro-pentofuranonucleosides were studied by Tyrsted et al. (*Biochim. Biophys. Acta* (1968), 155(2), 619-22) and Bloch, et al. (*J. Med. Chem.* (1967), 10(5), 908-12).

β -L-2'-deoxythymidine (β -L-dT) is known in the art to inhibit herpes simplex virus type 1 (HSV-1) thymidine kinase (TK). Iotti et al., WO 92/08727, teaches that β -L-dT selectively inhibits the phosphorylation of D-thymidine by HSV-1 TK, but not by human TK. Spaldari et al. reported that L-thymidine is phosphorylated by herpes simplex virus type 1 thymidine kinase and inhibits viral growth, *J. Med. Chem.* (1992), 35(22), 4214-20.

In light of the fact that hepatitis B virus has reached epidemic levels worldwide, and has severe and often tragic effects on the infected patient, there remains a strong need to provide new effective pharmaceutical agents to treat humans infected with the virus that have low toxicity to the host.

Therefore, it is an object of the present invention to provide new methods and compositions for the treatment of human patients or other hosts infected with hepatitis B virus.


Summary of the Invention

5

A method for the treatment of hepatitis B infection in humans and other host animals is disclosed that includes administering an effective amount of a biologically active 2'-deoxy- β -L-erythro-pentofuranonucleoside (referred to alternatively herein as a β -L-d-nucleoside or a β -L-2'-d-nucleoside) or a pharmaceutically acceptable salt or prodrug thereof, administered either alone or in combination, optionally in a pharmaceutically acceptable carrier. The term 2'-deoxy, as used in this specification, refers to a nucleoside that has no substituent in the 2'-position.

The disclosed 2'-deoxy- β -L-erythro-pentofuranonucleosides, or pharmaceutically acceptable prodrugs or salts or pharmaceutically acceptable formulations containing these compounds are useful in the prevention and treatment of hepatitis B infections and other related conditions such as anti-HBV antibody positive and HBV-positive conditions, chronic liver inflammation caused by HBV, cirrhosis, acute hepatitis, fulminant hepatitis, chronic persistent hepatitis, and fatigue. These compounds or formulations can also be used prophylactically to prevent or retard the progression of clinical illness in individuals who are anti-HBV antibody or HBV-antigen positive or who have been exposed to HBV.


In one embodiment of the present invention, the 2'-deoxy- β -L-erythro-pentofuranonucleoside derivative is a compound of the formula:



wherein R is selected from the group consisting of H, straight chained, branched or cyclic alkyl, CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, amino acid residue, mono, di, or triphosphate, or a phosphate

derivative; and BASE is a purine or pyrimidine base which may optionally be substituted.


In another embodiment, the 2'-deoxy- β -L-erythro-pentofuranonucleoside derivative is β -L-2'-deoxyadenosine or a pharmaceutically acceptable salt or prodrug thereof, of the formula:



5

wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

In another embodiment, the 2'-deoxy- β -L-erythro-pentofuranonucleoside derivative is β -L-2'-deoxycytidine or pharmaceutically acceptable salt or prodrug thereof of the formula:




10

wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).


In another embodiment, the 2'-deoxy- β -L-erythro-pentofuranonucleoside derivative is β -L-2'-deoxyuridine or pharmaceutically acceptable salt or prodrug thereof of the formula:

15




wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

In another embodiment, the 2'-deoxy-β-L-erythro-pentofuranonucleoside derivative is β-L-2'-deoxyguanosine or pharmaceutically acceptable salt or prodrug thereof of the formula:




wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

In another embodiment, the 2'-deoxy-β-L-erythro-pentofuranonucleoside derivative is β-L-2'-deoxyinosine or pharmaceutically acceptable salt or prodrug thereof of the formula:



wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

In another embodiment, the 2'-deoxy-β-L-erythro-pentofuranonucleoside derivative is β-L-thymidine or a pharmaceutically acceptable salt or prodrug thereof of the formula:



wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

In another embodiment, the 2'-deoxy-β-L-erythro-pentofuranonucleoside is administered in alternation or combination with one or more other 2'-deoxy-β-L-erythro-pentofuranonucleosides or one or more other compounds which exhibit activity against hepatitis B virus. In general, during alternation therapy, an effective dosage of each agent is administered serially, whereas in combination therapy, an effective dosage of two or more agents are administered together. The dosages will depend on absorption, inactivation, and excretion rates of the drug as well as other factors known to those of skill in the art. It is to be noted that dosage values will also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens and schedules should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions.

In another embodiment, the invention includes a method for the treatment of humans infected with HBV that includes administering an HBV treatment amount of a prodrug of the disclosed 2'-deoxy-β-L-erythro-pentofuranonucleoside derivatives. A prodrug, as used herein, refers to a compound that is converted into the nucleoside on administration in vivo.

Nonlimiting examples include pharmaceutically acceptable salt (alternatively referred to as "physiologically acceptable salts"), the 5' and N⁴ (cytidine) or N⁶ (adenosine) acylated or alkylated derivatives of the active compound, or the 5'-phospholipid or 5'-ether lipids of the active compound.

Brief Description of the Figure

Figure 1 illustrates a general process for obtaining β -L-erythro-pentafuranonucleosides (β -L-dN) using L-ribose or L-xylose as a starting material.

Figure 2 is a graph which illustrates the metabolism of L-dA, L-dC, and L-dT in human Hep G2 cells in terms of accumulation and decay. The cells were incubated with 10 μ M of compound.

Figure 3 is a graph which illustrates the antiviral effect of β -L-dA, β -L-dT and β -L-dC in the woodchuck chronic hepatitis model.

Detailed Description of the Invention

As used herein, the term “substantially in the form of a single isomer” or “in isolated form” refers to a 2'-deoxy- β -L-erythro-pentofuranonucleoside that is at least approximately 95% in the designated stereoconfiguration. In a preferred embodiment, the active compound is administered in at least this level of purity to the host in need of therapy.

As used herein, the term hepatitis B and related conditions refers to hepatitis B and related conditions such as anti-HBV antibody positive and HBV-positive conditions, chronic liver inflammation caused by HBV, cirrhosis, acute hepatitis, fulminant hepatitis, chronic persistent hepatitis, and fatigue. The method of the present invention includes the use of 2'-deoxy- β -L-erythro-pentofuranonucleoside derivatives prophylactically to prevent or retard the progression of clinical illness in individuals who are anti-HBV antibody or HBV-antigen positive or who have been exposed to HBV.

As used herein, the term alkyl, unless otherwise specified, refers to a saturated straight, branched, or cyclic, primary, secondary, or tertiary hydrocarbon, typically of C_1 to C_{18} , preferably C_1 to C_6 and specifically includes but is not limited to methyl, ethyl, propyl, butyl, pentyl, hexyl, isopropyl, isobutyl, sec-butyl, t-butyl, isopentyl, amyl, t-pentyl, cyclopentyl, and cyclohexyl.

As used herein, the term acyl refers to moiety of the formula $-C(O)R'$, wherein R' is alkyl; aryl, alkaryl, aralkyl, heteroaromatic, alkoxyalkyl including methoxymethyl; arylalkyl

including benzyl; aryloxyalkyl such as phenoxymethyl; aryl including phenyl optionally substituted with halogen, C₁ to C₄ alkyl or C₁ to C₄ alkoxy, or the residue of an amino acid. The term acyl specifically includes but is not limited to acetyl, propionyl, butyryl, pentanoyl, 3-methylbutyryl, hydrogen succinate, 3-chlorobenzoate, benzoyl, acetyl, pivaloyl, mesylate, propionyl, valeryl, caproic, caprylic, capric, lauric, myristic, palmitic, stearic, and oleic.

As used herein, the term purine or pyrimidine base, includes, but is not limited to, 6-alkylpurine and N⁶-alkylpurines, N⁶-acylpurines, N⁶-benzylpurine, 6-halopurine, N⁶-vinylpurine, N⁶-acetylenic purine, N⁶-acyl purine, N⁶-hydroxyalkyl purine, N⁶-thioalkyl purine, N²-alkylpurines, N⁴-alkylpyrimidines, N⁴-acylpyrimidines, 4-benzylpyrimidine, N⁴-halopyrimidines, N⁴-acetylenic pyrimidines, 4-acyl and N⁴-acyl pyrimidines, 4-hydroxyalkyl pyrimidines, 4-thioalkyl pyrimidines, thymine, cytosine, 6-azapyrimidine, including 6-azacytosine, 2- and/or 4-mercaptopyrimidine, uracil, C⁵-alkylpyrimidines, C⁵-benzylpyrimidines, C⁵-halopyrimidines, C⁵-vinylpyrimidine, C⁵-acetylenic pyrimidine, C⁵-acyl pyrimidine, C⁵-hydroxyalkyl purine, C⁵-amidopyrimidine, C⁵-cyanopyrimidine, C⁵-nitropyrimidine, C⁵-aminopyrimidine, N²-alkylpurines, N²-alkyl-6-thiopurines, 5-azacytidinyl, 5-azauracilyl, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, and pyrazolopyrimidinyl. Functional oxygen and nitrogen groups on the base can be protected as necessary or desired. Suitable protecting groups are well known to those skilled in the art, and include trimethylsilyl, dimethylhexylsilyl, *t*-butyldimethylsilyl, and *t*-butyldiphenylsilyl, trityl, alkyl groups, acyl groups such as acetyl and propionyl, methanesulfonyl, and *p*-toluenesulfonyl.

The term biologically active nucleoside, as used herein, refers to a nucleoside which exhibits an EC₅₀ of 15 micromolar or less when tested in 2.2.15 cells transfected with the hepatitis virion.

Preferred bases include cytosine, 5-fluorocytosine, 5-bromocytosine, 5-iodocytosine, uracil, 5-fluorouracil, 5-bromouracil, 5-iodouracil, 5-methyluracil, thymine, adenine, guanine, inosine, xanthine, 2,6-diaminopurine, 6-aminopurine, 6-chloropurine and 2,6-dichloropurine, 6-bromopurine, 2,6-dibromopurine, 6-iodopurine, 2,6-di-iodopurine, 5-bromovinylcytosine, 5-bromovinyluracil, 5-bromoethenylcytosine, 5-bromoethenyluracil, 5-trifluoromethylcytosine, 5-trifluoromethyluracil.

The 2'-deoxy- β -L-erythro-pentofuranonucleoside can be provided as a 5' phospholipid or a 5'-ether lipid, as disclosed in the following references, which are incorporated by reference herein: Kucera, L.S., N. Iyer, E. Leake, A. Raben, Modest E.J., D. L.W., and C. Piantadosi. 1990. Novel membrane-interactive ether lipid analogs that inhibit infectious HIV-1 production and induce defective virus formation. *AIDS Res Hum Retroviruses*. **6**:491-501; Piantadosi, C., J. Marasco C.J., S.L. morris-Natschke, K.L. Meyer, F. Gumus, J.R. Surles, K.S. Ishaq, L.S. Kucera, N. Iyer, C.A. Wallen, S. Piantadosi, and E.J. Modest. 1991-Synthesis and evaluation of novel ether lipid nucleoside conjugates for anti-HIV activity. *J Med Chem*. **34**:1408-1414; Hostetler, K.Y., D.D. Richman, D.A. Carson, L.M. Stuhmiller, G.M. T. van Wijk, and H. van den Bosch. 1992. Greatly enhanced inhibition of human immunodeficiency virus type 1 replication in CEM and HT4-6C cells by 31-deoxythymidine diphosphate dimyristoylglycerol, a lipid prodrug of 31-deoxythymidine. *Antimicrob Agents Chemother*. **36**:2025-2029; Hostetler, K.Y., L.M. Stuhmiller, H.B. Lenting, H. van den Bosch, and D.D. Richman. 1990. Synthesis and antiretroviral activity of phospholipid analogs of azidothymidine and other antiviral nucleosides. *J. Biol Chem*. **265**:6112-7.

The 2'-deoxy- β -L-erythro-pentofuranonucleoside can be converted into a pharmaceutically acceptable ester by reaction with an appropriate esterifying agent, for example, an acid halide or anhydride. The nucleoside or its pharmaceutically acceptable prodrug can be converted into a pharmaceutically acceptable salt thereof in a conventional manner, for example, by treatment with an appropriate base or acid. The ester or salt can be converted into the parent nucleoside, for example, by hydrolysis.

As used herein, the term pharmaceutically acceptable salts or complexes refers to salts or complexes of the 2'-deoxy- β -L-erythro-pentofuranonucleosides that retain the desired biological activity of the parent compound and exhibit minimal, if any, undesired toxicological effects. Nonlimiting examples of such salts are (a) acid addition salts formed with inorganic acids (for example, hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid, and the like), and salts formed with organic acids such as acetic acid, oxalic acid, tartaric acid, succinic acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmoic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acids, naphthalenedisulfonic acids, and polygalacturonic acid; (b) base addition salts formed with cations such as sodium, potassium, zinc, calcium, bismuth, barium,

magnesium, aluminum, copper, cobalt, nickel, cadmium, sodium, potassium, and the like, or with an organic cation formed from N,N-dibenzylethylene-diamine, ammonium, or ethylenediamine; or (c) combinations of (a) and (b); e.g., a zinc tannate salt or the like.

The term prodrug, as used herein, refers to a compound that is converted into the nucleoside on administration in vivo. Nonlimiting examples are pharmaceutically acceptable salts (alternatively referred to as “physiologically acceptable salts”), the 5' and N⁴ or N⁶ acylated or alkylated derivatives of the active compound, and the 5'-phospholipid and 5'-ether lipid derivatives of the active compound.

Modifications of the active compounds, specifically at the N⁴, N⁶ and 5'-O positions, can affect the bioavailability and rate of metabolism of the active species, thus providing control over the delivery of the active species.

A preferred embodiment of the present invention is a method for the treatment of HBV infections in humans or other host animals, that includes administering an effective amount of one or more of a 2'-deoxy-β-L-erythro-pentofuranonucleoside derivative selected from the group consisting of β-L-2'-deoxyadenosine, β-L-2'-deoxycytidine, β-L-2'-deoxyuridine, β-L-2'-guanosine, β-L-2'-deoxyinosine, and β-L-2'-deoxythymidine, or a physiologically acceptable prodrug thereof, including a phosphate, 5' and or N⁶ alkylated or acylated derivative, or a physiologically acceptable salt thereof, optionally in a pharmaceutically acceptable carrier. The compounds of this invention either possess anti-HBV activity, or are metabolized to a compound or compounds that exhibit anti-HBV activity. In a preferred embodiment, the 2'-deoxy-β-L-erythro-pentofuranonucleoside is administered substantially in the form of a single isomer, i.e., at least approximately 95% in the designated stereoconfiguration.

Nucleotide Prodrugs

Any of the nucleosides described herein can be administered as a stabilized nucleotide prodrug to increase the activity, bioavailability, stability or otherwise alter the properties of the nucleoside. A number of nucleotide prodrug ligands are known. In general, alkylation, acylation or other lipophilic modification of the mono, di or triphosphate of the nucleoside will increase the stability of the nucleotide. Examples of substituent groups that can replace one or more hydrogens on the phosphate moiety are alkyl, aryl, steroids, carbohydrates, including sugars,

1,2-diacylglycerol and alcohols. Many are described in R. Jones and N. Bischofberger, *Antiviral Research*, **27** (1995) 1-17. Any of these can be used in combination with the disclosed nucleosides to achieve a desired effect.

In one embodiment, the 2'-deoxy- β -L-erythro-pentofuranonucleoside is provided as 5'-hydroxyl lipophilic prodrug. Nonlimiting examples of U.S. patents that disclose suitable lipophilic substituents that can be covalently incorporated into the nucleoside, preferably at the 5'-OH position of the nucleoside or lipophilic preparations, include U.S. Patent Nos. 5,149,794 (Sep. 22, 1992, Yatvin et al.); 5,194,654 (Mar. 16, 1993, Hostetler et al.); 5,223,263 (June 29, 1993, Hostetler et al.); 5,256,641 (Oct. 26, 1993, Yatvin et al.); 5,411,947 (May 2, 1995, Hostetler et al.); 5,463,092 (Oct. 31, 1995, Hostetler et al.); 5,543,389 (Aug. 6, 1996, Yatvin et al.); 5,543,390 (Aug. 6, 1996, Yatvin et al.); 5,543,391 (Aug. 6, 1996, Yatvin et al.); and 5,554,728 (Sep. 10, 1996; Basava et al.), all of which are incorporated herein by reference.

Foreign patent applications that disclose lipophilic substituents that can be attached to the 2'-deoxy- β -L-erythro-pentofuranonucleoside derivative of the present invention, or lipophilic preparations, include WO 89/02733, WO 90/00555, WO 91/16920, WO 91/18914, WO 93/00910, WO 94/26273, WO 96/15132, EP 0 350 287, EP 93917054.4, and WO 91/19721.

Additional nonlimiting examples of 2'-deoxy- β -L-erythro-pentofuranonucleosides are those that contain substituents as described in the following publications. These derivatized 2'-deoxy- β -L-erythro-pentofuranonucleosides can be used for the indications described in the text or otherwise as antiviral agents, including as anti-HBV agents. Ho, D.H.W. (1973) Distribution of kinase and deaminase of 1 β -D-arabinofuranosylcytosine in tissues of man and mouse. *Cancer Res.* **33**, 2816-2820; Holy, A. (1993) Isopolar phosphorous-modified nucleotide analogues. In: De Clercq (Ed.), *Advances in Antiviral Drug Design*, Vol. I, JAI Press, pp. 179-231; Hong, C.I., Nechaev, A., and West, C.R. (1979a) Synthesis and antitumor activity of 1- β -D-arabinofuranosylcytosine conjugates of cortisol and cortisone. *Biochem. Biophys. Res. Commun.* **88**, 1223-1229; Hong, C.I., Nechaev, A., Kirisits, A.J. Buchheit, D.J. and West, C.R. (1980) Nucleoside conjugates as potential antitumor agents. 3. Synthesis and antitumor activity of 1-(β -D-arabinofuranosyl)cytosine conjugates of corticosteroids and selected lipophilic alcohols. *J. Med. Chem.* **28**, 171-177; Hostetler, K.Y., Stuhmiller, L.M., Lenting, H.B.M. van den Bosch, H. and Richman, D.D. (1990) Synthesis and antiretroviral activity of phospholipid analogs of

azidothymidine and other antiviral nucleosides. *J. Biol. Chem.* **265**, 6112-6117; Hostetler, K.Y., Carson, D.A. and Richman, D.D. (1991); Phosphatidylazidothymidine: mechanism of antiretroviral action in CEM cells. *J. Biol. Chem.* **266**, 11714-11717; Hostetler, K.Y., Korba, B. Sridhar, C., Gardener, M. (1994a) Antiviral activity of phosphatidyl-dideoxycytidine in hepatitis B-infected cells and enhanced hepatic uptake in mice. *Antiviral Res.* **24**, 59-67; Hostetler, K.Y., Richman, D.D., Sridhar, C.N. Felgner, P.L., Felgner, J., Ricci, J., Gardener, M.F. Selleseth, D.W. and Ellis, M.N. (1994b) Phosphatidylazidothymidine and phosphatidyl-ddC: Assessment of uptake in mouse lymphoid tissues and antiviral activities in human immunodeficiency virus-infected cells and in raucher leukemia virus-infected mice. *Antimicrobial Agents Chemother.* **38**, 2792-2797; Hunston, R.N., Jones, A.A. McGuigan, C., Walker, R.T., Balzarini, J., and De Clercq, E. (1984) Synthesis and biological properties of some cyclic phosphotriesters derived from 2'-deoxy-5-fluorouridine. *J. Med. Chem.* **27**, 440-444; Ji, Y.H., Moog, C., Schmitt, G., Bischoff, P. and Luu, B. (1990); Monophosphoric acid diesters of 7 β -hydroxycholesterol and of pyrimidine nucleosides as potential antitumor agents: synthesis and preliminary evaluation of antitumor activity. *J. Med. Chem.* **33**, 2264-2270; Jones, A.S., McGuigan, C., Walker, R.T., Balzarini, J. and DeClercq, E. (1984) Synthesis, properties, and biological activity of some nucleoside cyclic phosphoramidates. *J. Chem. Soc. Perkin Trans. I*, 1471-1474; Juodka, B.A. and Smart, J. (1974) Synthesis of ditribonucleoside a(P \rightarrow N) amino acid derivatives. *Coll. Czech. Chem. Comm.* **39**, 363-968; Kataoka, S., Imai, J., Yamaji, N., Kato, M., Saito, M., Kawada, T. and Imai, S. (1989) Alkylated cAMP derivatives; selective synthesis and biological activities. *Nucleic Acids Res. Sym. Ser.*, **21**, 1-2; Kataoka, S., Uchida, R. and Yamaji, N. (1991) A convenient synthesis of adenosine 3',5' cyclic phosphate (cAMP) benzyl and methyl triesters. *Heterocycles* **32**, 1351-1356; Kinchington, D., Harvey, J.J., O'Connor, T.J., Jones, B.C.N.M., Devine, K.G., Taylor-Robinson, D., Jeffries, D.J. and McGuigan, C. (1992) Comparison of antiviral effects of zidovudine phosphoramidate and phosphorodiamidate derivatives against HIV and MuLV *in vitro*. *Antiviral Chem. Chemother.* **3**, 107-112; Kodama, K., Morozumi, M., Saitoh, K.I., Kuninaka, H., Yoshino, H. and Saneyoshi, M. (1989) Antitumor activity and pharmacology of 1- β -D-arabinofuranosylcytosine -5'-stearylphosphate; an orally active derivative of 1- β -D-arabinofuranosylcytosine. *Jpn. J. Cancer Res.* **80**, 679-685; Korthy, M. and Engels, J. (1979) The effects of adenosine- and guanosine 3',5'-phosphoric and acid benzyl

esters on guinea-pig ventricular myocardium. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 310, 103-111; Kumar, A., Goe, P.L., Jones, A.S. Walker, R.T. Balzarini, J. and De Clercq, E. (1990) Synthesis and biological evaluation of some cyclic phosphoramidate nucleoside derivatives. *J. Med. Chem.* **33**, 2368-2375; LeBec, C., and Huynh-Dinh, T. (1991) Synthesis of lipophilic phosphate triester derivatives of 5-fluorouridine and arabinocytidine as anticancer prodrugs. *Tetrahedron Lett.* **32**, 6553-6556; Lichtenstein, J., Barner, H.D. and Cohen, S.S. (1960) The metabolism of exogenously supplied nucleotides by *Escherichia coli.*, *J. Biol. Chem.* **235**, 457-465; Luchty, J., Von Daeniken, A., Friederich, J. Manthey, B., Zweifel, J., Schlatter, C. and Benn, M.H. (1981) Synthesis and toxicological properties of three naturally occurring cyanoepithioalkanes. *Mitt. Geg. Lebensmittelunters. Hyg.* **72**, 131-133 (*Chem. Abstr.* 95, 127093); McGuigan, C. Tollerfield, S.M. and Riley, P.A. (1989) Synthesis and biological evaluation of some phosphate triester derivatives of the anti-viral drug Ara. *Nucleic Acids Res.* **17**, 6065-6075; McGuigan, C., Devine, K.G., O'Connor, T.J., Galpin, S.A., Jeffries, D.J. and Kinchington, D. (1990a) Synthesis and evaluation of some novel phosphoramidate derivatives of 3'-azido-3'-deoxythymidine (AZT) as anti-HIV compounds. *Antiviral Chem. Chemother.* **1**, 107-113; McGuigan, C., O'Connor, T.J., Nicholls, S.R. Nickson, C. and Kinchington, D. (1990b) Synthesis and anti-HIV activity of some novel substituted dialkyl phosphate derivatives of AZT and ddCyd. *Antiviral Chem. Chemother.* **1**, 355-360; McGuigan, C., Nicholls, S.R., O'Connor, T.J., and Kinchington, D. (1990c) Synthesis of some novel dialkyl phosphate derivative of 3'-modified nucleosides as potential anti-AIDS drugs. *Antiviral Chem. Chemother.* **1**, 25-33; McGuigan, C., Devine, K.G., O'Connor, T.J., and Kinchington, D. (1991) Synthesis and anti-HIV activity of some haloalkyl phosphoramidate derivatives of 3'-azido-3'-deoxythymidine (AZT); potent activity of the trichloroethyl methoxyalaninyl compound. *Antiviral Res.* **15**, 255-263; McGuigan, C., Pathirana, R.N., Mahmood, N., Devine, K.G. and Hay, A.J. (1992) Aryl phosphate derivatives of AZT retain activity against HIV-1 in cell lines which are resistant to the action of AZT. *Antiviral Res.* **17**, 311-321; McGuigan, C., Pathirana, R.N., Choi, S.M., Kinchington, D. and O'Connor, T.J. (1993a) Phosphoramidate derivatives of AZT as inhibitors of HIV; studies on the carboxyl terminus. *Antiviral Chem. Chemother.* **4**, 97-101; McGuigan, C., Pathirana, R.N., Balzarini, J. and De Clercq, E. (1993b) Intracellular

delivery of bioactive AZT nucleotides by aryl phosphate derivatives of AZT. *J. Med. Chem.* **36**, 1048-1052.

The question of chair-twist equilibria for the phosphate rings of nucleoside cyclic 3',5'-monophosphates. ¹HNMR and x-ray crystallographic study of the diastereomers of thymidine phenyl cyclic 3',5'-monophosphate. *J. Am. Chem. Soc.* **109**, 4058-4064; Nerbonne, J.M., Richard, S., Nargeot, J. and Lester, H.A. (1984) New photoactivatable cyclic nucleotides produce intracellular jumps in cyclic AMP and cyclic GMP concentrations. *Nature* **301**, 74-76; Neumann, J.M., Hervé, M., Debouzy, J.C., Guerra, F.I., Gouyette, C., Dupraz, B. and Huynh-Dinh, T. (1989) Synthesis and transmembrane transport studies by NMR of a glucosyl phospholipid of thymidine. *J. Am. Chem. Soc.* **111**, 4270-4277; Ohno, R., Tatsumi, N., Hirano, M., Imai, K., Mizoguchi, H., Nakamura, T., Kosaka, M., Takatuski, K., Yamaya, T., Toyama, K., Yoshida, T., Masaoka, T., Hashimoto, S., Ohshima, T., Kimura, I., Yamada, K. and Kimura, J. (1991) Treatment of myelodysplastic syndromes with orally administered 1-β-D-rabinofuranosylcytosine -5'-stearylphosphate. *Oncology* **48**, 451-455.

Palomino, E., Kessle, D. and Horwitz, J.P. (1989) A dihydropyridine carrier system for sustained delivery of 2',3'-dideoxynucleosides to the brain. *J. Med. Chem.* **32**, 622-625; Perkins, R.M., Barney, S., Wittrock, R., Clark, P.H., Levin, R., Lambert, D.M., Petteway, S.R., Serafinowska, H.T., Bailey, S.M., Jackson, S., Harnden, M.R., Ashton, R., Sutton, D., Harvey, J.J. and Brown, A.G. (1993) Activity of BRL47923 and its oral prodrug, SB203657A against a rauscher murine leukemia virus infection in mice. *Antiviral Res.* **20** (Suppl. I), 84; Piantadosi, C., Marasco, C.J., Jr., Morris-Natschke, S.L., Meyer, K.L., Gumus, F., Surles, J.R., Ishaq, K.S., Kucera, L.S., Iyer, N., Wallen, C.A., Piantadosi, S. and Modest, E.J. (1991) Synthesis and evaluation of novel ether lipid nucleoside conjugates for anti-HIV-1 activity. *J. Med. Chem.* **34**, 1408-1414; Pompon, A., Lefebvre, I., Imbach, J.L., Kahn, S. and Farquhar, D. (1994) Decomposition pathways of the mono- and bis(pivaloyloxymethyl) esters of azidothymidine-5'-monophosphate in cell extract and in tissue culture medium; an application of the 'on-line ISRP-cleaning' HPLC technique. *Antiviral Chem. Chemother.* **5**, 91-98; Postemark, T. (1974) Cyclic AMP and cyclic GMP. *Annu. Rev. Pharmacol.* **14**, 23-33; Prisbe, E.J., Martin, J.C.M., McGee, D.P.C., Barker, M.F., Smee, D.F., Duke, A.E., Matthews, T.R. and Verheyden, J.P.J. (1986) Synthesis and antiherpes virus activity of phosphate and phosphonate derivatives of 9-[(1,3-

dihydroxy-2-propoxy)methyl] guanine. *J. Med. Chem.* 29, 671-675; Puech, F., Gosselin, G., Lefebvre, I., Pompon, A., Aubertin, A.M. Dirn, A. and Imbach, J.L. (1993) Intracellular delivery of nucleoside monophosphate through a reductase-mediated activation process. *Antiviral Res.* 22, 155-174; Pugaeva, V.P., Klochkeva, S.I., Mashbits, F.D. and Eizengart, R.S. (1969). Robins, R.K. (1984) The potential of nucleotide analogs as inhibitors of retroviruses and tumors. *Pharm. Res.* 11-18; Rosowsky, A., Kim, S.H., Ross and J. Wick, M.M. (1982) Lipophilic 5'-(alkylphosphate) esters of 1- β -D-arabinofuranosylcytosine and its N^4 -acyl and 2,2'-anhydro-3'-O-acyl derivatives as potential prodrugs. *J. Med. Chem.* 25, 171-178; Ross, W. (1961) Increased sensitivity of the walker turnout towards aromatic nitrogen mustards carrying basic side chains following glucose pretreatment. *Biochem. Pharm.* 8, 235-240; Ryu, E.K., Ross, R.J. Matsushita, T., MacCoss, M., Hong, C.I. and West, C.R. (1982). Phospholipid-nucleoside conjugates. 3. Synthesis and preliminary biological evaluation of 1- β -D-arabinofuranosylcytosine 5'diphosphate[-], 2-diacylglycerols. *J. Med. Chem.* 25, 1322-1329; Saffhill, R. and Hume, W.J. (1986) The degradation of 5-iododeoxyuridine and 5-bromodeoxyuridine by serum from different sources and its consequences for the use of these compounds for incorporation into DNA. *Chem. Biol. Interact.* 57, 347-355; Saneyoshi, M., Morozumi, M., Kodama, K., Machida, J., Kuninaka, A. and Yoshino, H. (1980) Synthetic nucleosides and nucleotides. XVI. Synthesis and biological evaluations of a series of 1- β -D-arabinofuranosylcytosine 5'-alkyl or arylphosphates. *Chem. Pharm. Bull.* 28, 2915-2923; Sastry, J.K., Nehete, P.N., Khan, S., Nowak, B.J., Plunkett, W., Arlinghaus, R.B. and Farquhar, D. (1992) Membrane-permeable dideoxyuridine 5'-monophosphate analogue inhibits human immunodeficiency virus infection. *Mol. Pharmacol.* 41, 441-445; Shaw, J.P., Jones, R.J. Arimilli, M.N., Louie, M.S., Lee, W.A. and Cundy, K.C. (1994) Oral bioavailability of PMEA from PMEA prodrugs in male Sprague-Dawley rats. 9th Annual AAPS Meeting. San Diego, CA (Abstract). Shuto, S., Ueda, S., Imamura, S., Fukukawa, K. Matsuda, A. and Ueda, T. (1987) A facile one-step synthesis of 5'-phosphatidyl nucleosides by an enzymatic two-phase reaction. *Tetrahedron Lett.* 28, 199-202; Shuto, S., Itoh, H., Ueda, S., Imamura, S., Kukukawa, K., Tsujino, M., Matsuda, A. and Ueda, T. (1988) A facile enzymatic synthesis of 5'-(3-sn-phosphatidyl)nucleosides and their antileukemic activities. *Chem. Pharm. Bull.* 36, 209-217. One preferred phosphate prodrug group is the S-acyl-2-thioethyl group, also referred to as "SATE."

Combination or Alternation Therapy

It has been recognized that drug-resistant variants of HBV can emerge after prolonged treatment with an antiviral agent. Drug resistance most typically occurs by mutation of a gene that encodes for an enzyme used in the viral life cycle, and most typically in the case of HBV, DNA polymerase. Recently, it has been demonstrated that the efficacy of a drug against HBV infection can be prolonged, augmented, or restored by administering the compound in combination or alternation with a second, and perhaps third, antiviral compound that induces a different mutation from that caused by the principle drug. Alternatively, the pharmacokinetics, biodistribution, or other parameter of the drug can be altered by such combination or alternation therapy. In general, combination therapy is typically preferred over alternation therapy because it induces multiple simultaneous stresses on the virus.

The anti-hepatitis B viral activity of β -L-2'-dA, β -L-2'-dC, β -L-2'-dU, β -L-2'-dG, β -L-2'-dT, β -L-dI, or other β -L-2'-nucleosides provided herein, or the prodrugs, phosphates, or salts of these compounds, can be enhanced by administering two or more of these nucleosides in combination or alternation. Alternatively, for example, one or more of β -L-2'-dA, β -L-2'-dC, β -L-2'-dU, β -L-2'-dG, β -L-2'-dT, β -L-dI, or other β -L-2'-nucleosides provided herein can be administered in combination or alternation with 3TC, FTC, L-FMAU, DAPD, famciclovir, penciclovir, BMS-200475, bis pom PMEA (adefovir, dipivoxil); lobucavir, ganciclovir, or ribavirin.

In any of the embodiments described herein, if the β -L-2'-nucleoside of the present invention is administered in combination or alternation with a second nucleoside or nonnucleoside reverse transcriptase inhibitor that is phosphorylated to an active form, it is preferred that a second compound be phosphorylated by an enzyme that is different from that which phosphorylates the selected β -L-2'-nucleoside of the present invention in vivo. Examples of kinase enzymes are thymidine kinase, cytosine kinase, guanosine kinase, adenosine kinase, deoxycytidine kinase, 5'-nucleotidase, and deoxyguanosine kinase.

Preparation of the Active Compounds

The 2'-deoxy- β -L-erythro-pentofuranonucleoside derivatives of the present invention are known in the art and can be prepared according to the method disclosed by Holy, *Collect. Czech. Chem. Commun.* (1972), 37(12), 4072-87 and *Mol. Phys.* (1967), 3(4), 386-95.

A general process for obtaining β -L-erythro-pentafuranonucleosides (β -L-dN) is shown in Figure 1, using L-ribose or L-xylose as a starting material.


Mono, di, and triphosphate derivatives of the active nucleosides can be prepared as described according to published methods. The monophosphate can be prepared according to the procedure of Imai et al., *J. Org. Chem.*, 34(6), 1547-1550 (June 1969). The diphosphate can be prepared according to the procedure of Davisson et al., *J. Org. Chem.*, 52(9), 1794-1801 (1987). The triphosphate can be prepared according to the procedure of Hoard et al., *J. Am. Chem. Soc.*, 87(8), 1785-1788 (1965).

Experimental Protocols

Melting points were determined in open capillary tubes on a Gallenkamp MFB-595-010 M apparatus and are uncorrected. The UV absorption spectra were recorded on an Uvikon 931 (KONTRON) spectrophotometer in ethanol. ^1H -NMR spectra were run at room temperature in $\text{DMSO}-d_6$ with a Bruker AC 250 or 400 spectrometer. Chemical shifts are given in ppm, $\text{DMSO}-d_5$ being set at 2.49 ppm as reference. Deuterium exchange, decoupling experiments or 2D-COSY were performed in order to confirm proton assignments. Signal multiplicities are represented by s (singlet), d (doublet), dd (doublet of doublets), t (triplet), q (quadruplet), br (broad), m (multiplet). All J-values are in Hz. FAB mass spectra were recorded in the positive- (FAB>0) or negative- (FAB<0) ion mode on a JEOL DX 300 mass spectrometer. The matrix was 3-nitrobenzyl alcohol (NBA) or a mixture (50:50, v/v) of glycerol and thioglycerol (GT). Specific rotations were measured on a Perkin-Elmer 241 spectropolarimeter (path length 1 cm) and are given in units of $10^{-1} \text{ deg cm}^2 \text{ g}^{-1}$. Elemental analysis were carried out by the "Service de Microanalyses du CNRS, Division de Vernaison" (France). Analyses indicated by the symbols of the elements or functions were within $\pm 0.4\%$ of theoretical values. Thin layer chromatography was performed on precoated aluminium sheets of Silica Gel 60 F₂₅₄ (Merck, Art. 5554), visualisation of products being accomplished by UV absorbency followed by charring with 10%

ethanolic sulfuric acid and heating. Column chromatography was carried out on Silica Gel 60 (Merck, Art. 9385) at atmospheric pressure.

Example 1 Stereospecific Synthesis of 2'-Deoxy-β-L-Adenosine



9-(3,5-Di-*O*-benzoyl- β -L-xylofuranosyl)adenine (**3**)

A solution of 9-(2-*O*-acetyl-3,5-di-*O*-benzoyl- β -L-xylofuranosyl)adenine **2** [Ref.: Gosselin, G.; Bergogne, M.-C.; Imbach, J.-L., "Synthesis and Antiviral Evaluation of β -L-Xylofuranosyl Nucleosides of the Five Naturally Occurring Nucleic Acid Bases", *Journal of Heterocyclic Chemistry*, 1993, **30** (Oct.-Nov.), 1229-1233] (8.30 g, 16.05 mmol) and hydrazine hydrate 98% (234 mL, 48.5 mmol) in a mixture of pyridine / glacial acetic acid (4/1, v/v, 170 mL) was stirred at room temperature for 22 h. The reaction was quenched by adding acetone (40 mL) and stirring was continued for one additional hour. The reaction mixture was reduced to one half of its volume, diluted with water (250 mL) and extracted with chloroform (2 x 150 mL). The organic layer was washed successively with an aqueous saturated solution of NaHCO₃ (3 x 100 mL) and water (3 x 100 mL), dried, filtered, concentrated and co-evaporated with toluene and methanol. The residue was purified by silica gel column chromatography (0-3% MeOH in dichloromethane) to give **3** (5.2 g, 68%) precipitated from diisopropyl ether : ¹H NMR (DMSO-*d*₆) : δ 4.5-4.9 (m, 4H, H-2', H-4', H-5' and H-5''), 5.64 (t, 1H, H-3', J_{2',3'} = J_{3',4'} = 3.5 Hz), 6.3 (br s, 1H, OH-2'), 6.45 (d, 1H, H-1', J_{1',2'} = 4.6 Hz), 7.3 (br s, 2H, NH₂-6), 7.4-7.9 (m, 10H, 2 benzoyls), 8.07 and 8.34 (2s, 2H, H-2 and H-8); ms : matrix G/T, (FAB⁺) m/z 476 [M+H]⁺, 136 [BH₂]⁺, (FAB⁻) m/z 474 [M-H]⁻, 134 [B]⁻; UV (95% ethanol) : λ_{max} 257 nm (ϵ 16400), 230 nm (ϵ 29300), λ_{min} 246 nm (ϵ 14800); $[\alpha]_{\text{D}}^{20}$ = - 64 (c 1.07, CHCl₃). Anal. Calcd for C₂₄H₂₁N₅O₄ (M = 475.45) : C, 60.43; H, 4.45; N, 14.73. Found : C, 60.41; H, 4.68; N, 14.27.

9-(3,5-Di-*O*-benzoyl-2-deoxy- β -L-*threo*-pentofuranosyl)adenine (**4**).

To a solution of compound **3** (1.00 g, 2.11 mmol) in dry acetonitrile (65 mL) were added 4-(dimethylamino)pyridine (0.77 g, 6.32 mmol) and phenoxythiocarbonyl chloride (0.44 mL, 3.16 mmol). The mixture was stirred at room temperature for 2 h. After concentration, the residue was dissolved in dichloromethane (50 mL) and washed successively with water (2 x 30 mL), aqueous solution of hydrochloric acid 0.5 N (30 mL) and water (3 x 30 mL). The organic layer was dried, filtered and concentrated to dryness. The crude thiocarbonylated intermediate was directly treated with tris-(trimethylsilyl)silane hydride (0.78 mL, 5.23 mmol) and α,α' -azoisobutyronitrile (AIBN, 0.112 g, 0.69 mmol) in dry dioxane (17 mL) at reflux for 2 h. The solvent was removed under vacuum and the residue was purified by silica gel column chromatography (0-5% MeOH in dichloromethane) to give pure **4** (0.93 g, 96%) as a foam : ¹H

NMR (DMSO- d_6) : δ 2.9-3.1 (m, 2H, H-2' and H-2''), 4.6 - 4.7 (m, 3H, H-4', H-5' and H-5''), 5.8 (br s, 1H, H-3'), 6.43 (dd, 1H, H-1', $J_{1',2'} = 3.1$ Hz, $J_{1',2''} = 7.6$ Hz), 7.3 (br s, 2H, NH₂-6), 7.4-7.9 (m, 10H, 2 benzoyls), 8.05 and 8.33 (2s, 2H, H-2 and H-8); ms : matrix G/T, (FAB⁺) m/z 460 [M+H]⁺, 325 [S]⁺, 136 [BH₂]⁺, (FAB⁻) m/z 458 [M-H]⁻, 134 [B]⁻; UV (95% ethanol) : λ_{\max} 261 nm (ϵ 14400), 231 nm (ϵ 26300), λ_{\min} 249 nm (ϵ 12000); $[\alpha]_D^{20} = -38$ (c 1.04, DMSO).

6-N-(4-Monomethoxytrityl)-9-(3,5-di-O-benzoyl-2-deoxy- β -L-threo-pentofuranosyl)adenine (5).

To a solution of compound 4 (0.88 g, 1.92 mmol) in dry pyridine (40 mL) was added 4-monomethoxytrityl chloride (1.18 g, 3.84 mmol). The mixture was stirred at 60°C for 24 h. After addition of methanol (5 mL), the solution was concentrated to dryness, the residue was dissolved in dichloromethane (50 mL) and washed successively with water (30 mL), aqueous saturated NaHCO₃ (30 mL) and water (30 mL). The organic layer was dried, filtered, concentrated and co-evaporated with toluene to give pure 5 (1.01 g, 72%) as a foam : ¹H NMR (CDCl₃) : δ 2.9-3.0 (m, 2H, H-2' and H-2''), 3.62 (s, 3H, OCH₃), 4.6-4.8 (m, 3H, H-4', H-5' and H-5''), 5.85 (pt, 1H, H-3'), 6.44 (dd, 1H, H-1', $J_{1',2'} = 3.1$ Hz, $J_{1',2''} = 7.3$ Hz), 6.9 (br s, 1H, NH-6), 6.7-6.8 and 7.2-7.4 (2m, 24H, 2 benzoyls and MMTr), 7.97 and 8.13 (2s, 2H, H-2 and H-8); ms : matrix G/T, (FAB⁺) m/z 732 [M+H]⁺, (FAB⁻) m/z 730 [M-H]⁻; UV (95% ethanol) : λ_{\max} 274 nm (ϵ 12100), 225 nm (ϵ 24200), λ_{\min} 250 nm (ϵ 5900); $[\alpha]_D^{20} = -16$ (c 1.12, DMSO).

6-N-(4-Monomethoxytrityl)-9-(2-deoxy- β -L-threo-pentofuranosyl)-adenine (6).

Compound 5 (0.95 g, 1.30 mmol) was treated with a solution (saturated at -10°C) of methanolic ammonia (40 mL), at room temperature overnight. After concentration, the residue was dissolved in dichloromethane (60 mL) and washed with water (30 mL). The aqueous layer was extracted twice with dichloromethane (10 mL). The combined organic layer was dried, filtered and concentrated. The residue was purified by silica gel column chromatography (0-5% MeOH in dichloromethane) to give pure 6 (0.67 g, 98%) as a foam : ¹H NMR (CDCl₃) : δ 2.6-2.9 (m, 2H, H-2' and H-2''), 3.5 (br s, 1H, OH-5'), 3.55 (s, 3H, OCH₃), 3.9-4.0 (m, 3H, H-4', H-5' and H-5''), 4.5-4.6 (m, 1H, H-3'), 6.03 (dd, 1H, H-1', $J_{1',2'} = 4.0$ Hz, $J_{1',2''} = 8.8$ Hz), 7.0 (br s, 1H, NH-6), 6.7-6.8 and 7.1-7.4 (2m, 14H, MMTr), 7.40 (d, 1H, OH-3', $J_{H,OH} = 10.6$ Hz), 7.80 and 7.99 (2s, 2H, H-2 and H-8); ms : matrix G/T, (FAB⁺) m/z 524 [M+H]⁺, 408 [BH₂]⁺, (FAB⁻)

m/z 1045 [2M-H]⁺, 522 [M-H]⁺, 406 [B]⁺; UV (95% ethanol) : λ_{\max} 275 nm (ϵ 12300), λ_{\min} 247 nm (ϵ 3600); $[\alpha]_D^{20} = +28$ (c 0.94, DMSO).

6-N-(4-Monomethoxytrityl)-9-(2-deoxy-5-O-(4-monomethoxytrityl)- β -L-threo-pentofuranosyl)adenine (7).

Compound **6** (0.62 g, 1.24 mmol) in dry pyridine (25 mL) was treated with 4-monomethoxytrityl chloride (0.46 g, 1.49 mmol) at room temperature for 16 h. After addition of methanol (5 mL), the mixture was concentrated to dryness. The residue was dissolved in dichloromethane (60 mL) and washed successively with water (40 mL), a saturated aqueous solution of NaHCO₃ (40 mL) and water (3 x 40 mL). The organic layer was dried, filtered, concentrated and co-evaporated with toluene and methanol. The residue was purified by silica gel column chromatography (0-10% MeOH in dichloromethane) to give **7** (0.71 g, 72%) as a foam : ¹H NMR (DMSO-*d*₆) : δ 2.21 (d, 1H, H-2' J_{2',2''} = 14.3 Hz), 2.6-2.7 (m, 1H, H-2''), 3.1-3.3 (2m, 2H, H-5' and H-5''), 3.64 and 3.65 (2s, 6H, 2 x OCH₃), 4.1-4.2 (m, 1H, H-4'), 4.2-4.3 (m, 1H, H-3'), 5.68 (d, 1H, OH-3', J_{H,OH} = 5.2 Hz), 6.24 (d, 1H, H-1', J_{1',2''} = 7.0 Hz), 6.7-6.8 and 7.1-7.3 (2m, 29H, 2 MMTr and NH-6), 7.83 and 8.21 (2s, 2H, H-2 and H-8); ms : matrix G/T, (FAB⁺) m/z 796 [M+H]⁺, 408 [BH₂]⁺, (FAB⁻) m/z 794 [M-H]⁻, 406 [B]⁻; UV (95% ethanol) : λ_{\max} 275 nm (ϵ 30900), λ_{\min} 246 nm (ϵ 12800); $[\alpha]_D^{20} = +14$ (c 1.03, DMSO).

6-N-(4-Monomethoxytrityl)-9-(3-O-benzoyl-2-deoxy-5-O-(4-mono-methoxytrityl)- β -L-erythro-pentofuranosyl)adenine (8).

A solution of diethylazodicarboxylate (0.38 mL, 2.49 mmol) in dry tetrahydrofuran (20 mL) was added dropwise to a cooled solution (0°C) of nucleoside **7** (0.66 g, 0.83 mmol), triphenylphosphine (0.66 g, 2.49 mmol) and benzoic acid (0.30 g, 2.49 mmol) in dry THF (20 mL). The mixture was stirred at room temperature for 18 h and methanol (1 mL) was added. The solvents were removed under reduced pressure and the crude material was purified by silica gel column chromatography (0-5% ethyl acetate in dichloromethane) to give compound **8** slightly contaminated by triphenylphosphine oxide.

6-N-(4-Monomethoxytrityl)-9-(2-deoxy-5-O-(4-monomethoxytrityl)- β -L-erythro-pentofuranosyl)adenine (9).

Compound **8** was treated by a solution (saturated at -10°C) of methanolic ammonia (20 mL), at room temperature for 24 h, then the reaction mixture was concentrated to dryness. The

residue was dissolved in dichloromethane (30 mL) and washed with water (20 mL). The aqueous layer was extracted by dichloromethane (2 x 20 mL) and the combined organic phase was dried, filtered and concentrated. Pure compound **9** (0.50 g, 76% from **7**) was obtained as a foam after purification by silica gel column chromatography (0-2% MeOH in dichloromethane) : ^1H NMR (DMSO- d_6) : δ 2.2-2.3 (m, 1H, H-2'), 2.8-2.9 (m, 1H, H-2''), 3.1-3.2 (m, 2H, H-5' and H-5''), 3.64 and 3.65 (2s, 6H, 2 x OCH₃), 3.97 (pq, 1H, H-4'), 4.4-4.5 (m, 1H, H-3'), 5.36 (d, 1H, OH-3', $J_{\text{H,OH}} = 4.5$ Hz), 6.34 (t, 1H, H-1', $J_{1',2'} = J_{1',2''} = 6.4$ Hz), 6.8-6.9 and 7.1-7.4 (2m, 29H, 2 MMTr and NH-6), 7.81 and 8.32 (2s, 2H, H-2 and H-8); ms : matrix G/T, (FAB⁺) m/z 796 [M+H]⁺, 408 [BH₂]⁺, (FAB⁻) m/z 794 [M-H]⁻, 406 [B]⁻; UV (95% ethanol) : λ_{max} 276 nm (ϵ 42600), λ_{min} 248 nm (ϵ 23300); $[\alpha]_{\text{D}}^{20} = +29$ (c 1.05, DMSO).

2'-Deoxy- β -L-adenosine (β -L-dA)


Compound **9** (0.44 g, 0.56 mmol) was treated with an aqueous solution of acetic acid 80% (17 mL) at room temperature for 5 h. The mixture was concentrated to dryness, the residue was dissolved in water (20 mL) and washed with diethyl ether (2 x 15 mL). The aqueous layer was concentrated and co-evaporated with toluene and methanol. The desired 2'-deoxy- β -L-adenosine (**β -L-dA**) (0.12 g, 83%) was obtained after purification by silica gel column chromatography (0-12% MeOH in dichloromethane) and filtration through a Millex HV-4 unit (0.45 μ , Millipore) : mp 193-194°C (crystallized from water) (Lit. 184-185°C for L-enantiomer [Ref.: Robins, M. J.; Khwaja, T. A.; Robins, R. K. *J. Org. Chem.* **1970**, *35*, 636-639] and 187-189°C for D-enantiomer [Ref.: Ness, R. K. in *Synthetic Procedures in Nucleic Acid Chemistry*; Zorbach, W.W., Tipson, R. S., Eds.; J. Wiley and sons : New York, **1968**; *Vol 1*, pp 183-187]; ^1H NMR (DMSO- d_6) : δ 2.2-2.3 and 2.6-2.7 (2m, 2H, H-2' and H-2''), 3.4-3.6 (2m, 2H, H-5' and H-5''), 3.86 (pq, 1H, H-4'), 4.3-4.4 (m, 1H, H-3'), 5.24 (t, 1H, OH-5', $J_{\text{H,OH}} = 5.8$ Hz), 5.30 (d, 1H, OH-3', $J_{\text{H,OH}} = 4.0$ Hz), 6.32 (dd, 1H, H-1', $J_{1',2'} = 6.2$ Hz, $J_{1',2''} = 7.8$ Hz), 7.3 (br s, 2H, NH₂-6), 8.11 and 8.32 (2s, 2H, H-2 and H-8); ms : matrix G/T, (FAB⁺) m/z 252 [M+H]⁺, 136 [BH₂]⁺, (FAB⁻) m/z 250 [M-H]⁻, 134 [B]⁻; UV (95% ethanol) : λ_{max} 258 nm (ϵ 14300), λ_{min} 226 nm (ϵ 2100); $[\alpha]_{\text{D}}^{20} = +25$ (c 1.03, H₂O), (Lit. $[\alpha]_{\text{D}}^{20} = +23$ (c 1.0, H₂O) for L-enantiomer [Ref.: Robins, M. J.; Khwaja, T. A.; Robins, R. K. *J. Org. Chem.* **1970**, *35*, 636-639] and $[\alpha]_{\text{D}}^{20} = -25$ (c 0.47, H₂O) for D-enantiomer [Ref.: Ness, R. K. in *Synthetic Procedures in Nucleic Acid Chemistry*; Zorbach, W.W., Tipson, R. S., Eds.; J. Wiley and sons : New York, **1968**; *Vol 1*, pp

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2423	2424	2425	2426	2427	2428	2429	2430	2431	2432	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

Example 2 Stereoselective Synthesis of 2'-Deoxy- β -L-Adenosine (β -L-dA)



Reaction 1:



Precursor: L-ribose (Cultor Science Food, CAS [24259-59-4], batch RIB9711013)

Reactants: Sulphuric acid 95-97% (Merck; ref 1.00731.1000); Benzoyl chloride (Fluka; ref 12930); Sodium sulfate (Prolabo; ref 28111.365)

Solvents: Methanol P.A. (Prolabo; ref 20847.295); Pyridine 99% (Acros; ref 131780025); Dichloromethane P.A. (Merck; ref 1.06050.6025); Acetic acid P.A. (carlo erba; ref 20104298); Acetic anhydride (Fluka; ref 45830); Ethanol 95 (Prolabo; ref 20823.293)

References: Recondo, E. F., and Rinderknecht, H., Eine neue, Einfache Synthese des 1-O-Acetyl-2,3,5-Tri-O-β-D-Ribofuranosides. *Helv. Chim. Acta*, 1171-1173 (1959).

A solution of L-ribose **140** (150 g, 1 mol) in methanol (2 liters) was treated with sulphuric acid (12 ml) and left at +4°C for 12hrs, and then neutralised with pyridine (180 ml). Evaporation gave an α,β mixture of methyl ribofuranosides **141** as a syrup. A solution of this anomeric mixture in pyridine (1.3 liters) was treated with benzoyl chloride (580 ml, 5 mol) with cooling and mechanical stirring. The solution was left at room temperature for 12 hrs and then poured on ice (about 10 liters) with continued stirring. The mixture (an oil in water) was filtered on a Cellite bed. The resulting oil on the cellite bed was washed with water (3x3 liters) and then dissolved with ethyl acetate (3 liters). The organic phase was washed with a 5% NaHCO_3 solution (2 liters) and water (2 liters), dried over sodium sulfate, filtered and evaporated to give 1-O-methyl-2,3,5-tri-O-benzoyl-α/β-L-ribofuranose **142** as a thick syrup. The oil was dissolved in acetic anhydride (560 ml) and acetic acid (240 ml). The solution was, after the dropwise addition of concentrated sulphuric acid (80 ml), kept in the cold (+4°C) under mechanical stirring for 10 hrs. The solution was then poured on ice (about 10 liters) under continued stirring. The mixture (oily compound in water) was filtered on a Cellite bed. The resulting gummy solid on the cellite bed was washed with water (3x3 liters) and then dissolved in dichloromethane (2,5

liters). The organic phase was washed with 5% NaHCO₃ (1 liter) and water (2x2 liters), dried over sodium sulfate, filtered and evaporated to give a gummy solid **143**, which was crystallized from ethanol 95 (yield 225 g, 44%).

Analyses for 1-O-acetyl-2,3,5-tri-O-benzoyl-β-L-ribofuranose 143 :


mp 129-130°C (EtOH 95) (lit.(1) mp 130-131°C)

¹H NMR (200 MHz, CDCl₃): δ 8.09-7.87 (m, 6H, H_{Arom}), 7.62-7.31 (m, 9H, H_{Arom}) 6.43 (s, 1H, H₁), 5.91 (dd, 1H, H₃, J_{3,4} 6.7 Hz; J_{3,2} 4.9 Hz), 5.79 (pd, 1H, H₂, J_{2,3} 4.9 Hz; J_{1,2} <1), 4.78 (m, 2H, H₄ and H₅), 4.51 (dd, 1H, H₅, J_{5,5'} 13.1 Hz, J_{5',4} 5.5 Hz), 2.00 (s, 3H, CH₃CO); (identical to commercial 1-O-acetyl-2,3,5-tri-O-benzoyl-β-D-ribofuranose)

Mass analysis (FAB+, GT) m/z 445 (M-OAc)+

Elemental analysis C₂₈H₂₄O₉ Calculated C 66.66 H 4.79; found C H

Reaction 2:



Precursor: Adenine (Pharma-Waldhof; ref 400134.001 lot 45276800)

Reactants: Stannic chloride fuming (Fluka; ref 96558); NH₃ / Methanol (methanol saturated with NH₃; see page 5); Sodium sulfate (Prolabo; ref 28111.365)

Solvents: Acetonitrile (Riedel-de Hean; ref 33019; distilled over CaH₂); Chloroform Pur (Acros; ref 22706463); Ethyl acetate Pur (Carlo erba; ref 528299)

References: Saneyoshi, M., and Satoh, E., Synthetic Nucleosides and Nucleotides. XIII. Stannic Chloride Catalyzed Ribosylation of Several 6-Substituted Purines. *Chem; Pharm. Bull.*, **27**, 2518-2521 (1979).; Nakayama, C., and Saneyoshi, M., Synthetic Nucleosides and Nucleotides. XX. Synthesis of Various 1-β-Xylofuranosyl-5-Alkyluracils and Related Nucleosides. *Nucleosides, Nucleotides*, **1**, 139-146 (1982).

Adenine (19.6 g, 144 mmol) was suspended in acetonitrile (400 ml) with 1-*O*-acetyl-2,3,5-tri-*O*-benzoyl- β -L-ribofuranose **143** (60 g, 119 mmol). To this suspension was added stannic chloride fuming (22 ml, 187 mmol). After 12 hrs, the reaction was concentrated to a small volume (about 100 ml), and sodium hydrogencarbonate (110 g) and water (120 ml) were added. The resulting white solid (tin salts) was extracted with hot chloroform (5x200 ml). The combined extracts were filtered on a cellite bed. The organic phase was washed with a NaHCO₃ 5% solution and water, dried over sodium sulfate, filtered and evaporated to give compound **144** (60 g, colorless foam). The foam was treated with methanol saturated with ammonia (220 ml) in sealed vessel at room temperature under stirring for 4 days. The solvent was evaporated off under reduced pressure and the resulting powder was suspended in ethyl acetate (400 ml) at reflux for 1 hr. After filtration, the powder was recrystallized from water (220 ml) to give L-adenosine **145** (24 g, crystals, 75%)


Analyses for β -L-adenosine:

mp 233-234°C (water) (lit.(4) mp 235°-238°C)

¹H NMR (200 MHz, DMSO-D₆): δ 8.34 and 8.12 (2s, 2H, H₂ and H₈), 7.37 (1s, 2H, NH₂), 5.86 (d, 1H, H_{1'}, J_{1',2'} 6.2 Hz), 5.43 (m, 2H, OH_{2'} and OH_{5'}), 5.19 (d, 1H, OH_{3'}, J 3.7 Hz), 4.60 (m, H_{2'}), 4.13 (m, 1H, H_{3'}), 3.94 (m, 1H, H_{4'}), 3.69-3.49 (m, 2H, H_{5'a} and H_{5'b}), (identical to commercial D-adenosine)

Mass analysis (FAB+, GT) m/z 268 (M+H)⁺, 136(BH₂)⁺

Reaction 3:



Reactants: 1,3-Dichloro-1,1,3,3-tetraisopropyldisiloxane (Fluka; ref 36520); Sodium sulfate (Prolabo; ref 28111.365)

Solvents: Pyridine 99% (Acros; ref 131780025); Ethyl acetate Pur (Carlo erba; ref 528299); Acetonitrile (Riedel-de Haen; ref 33019)

Reference : Robins, M.J., et al., Nucleic Acid Related Compounds. 42. A General Procedure for the Efficient Deoxygenation of Secondary Alcohols. Regiospecific and Stereoselective Conversion of Ribonucleosides to 2'-Deoxynucleosides. *J. Am. Chem. Soc.* **105**, 4059-4065 (1983).

To L-adenosine **145** (47,2 g, 177 mmol) suspended in pyridine (320 ml) was added 1,3-dichloro-1,1,3,3-tetraisopropyldisiloxane (63 ml, 201 mmol), and the mixture was stirred at room temperature for 12 hrs. Pyridine was evaporated and the residue was partitioned with ethyl acetate (1 liter) and a NaHCO₃ 5 % solution (600 ml). The organic phase was washed with a HCl 0.5N solution (2x500 ml) and water (500 ml), dried over sodium sulfate, filtered and evaporated to dryness. The resulting solid was crystallized from acetonitrile to give compound **146** (81 g, 90%).


Analyses 3',5'-O-(1,1,3,3-tetraisopropyl-1,3-disiloxanyl)-β-L-adenosine 146 :

mp 97-98°C (acetonitrile) (lit. (5) D enantiomer mp 98°C)

¹H NMR (200 MHz, CDCl₃): δ 8.28 and 7.95 (2s, 2H, H₂ and H₈), 5.96 (d, 1H, J_{1',2'} 1,1 Hz), 5.63 (s, 2H, NH₂), 5.10 (dd, 1H, H_{3'}, J_{3',4'} 7.6 Hz, J_{3',2'} 5.5 Hz), 4.57 (dd, 1H, H_{2'}, J_{2',1'} 1.2 Hz; J_{2',3'} 7.6 Hz), 4.15-3.99 (m, 3H, H_{4'}, H_{5'a} and H_{5'b}), 3.31 (sl, 1H, OH_{2'}), 1.06 (m, 28H, isopropyl protons)

Mass analysis (FAB-, GT) m/z 508 (M-H)⁻, 134 (B)⁻; (FAB+, GT) m/z 510 (m+H)⁺, 136 (BH₂)⁺

Reaction 4:



Reactants: Dimethylaminopyridine 99% (Acros; ref 1482702050);

Phenylchlorothionocarbonate 99% (Acros; ref 215490050); Tris(trimethylsilyl)silane "TTMSS" (Fluka; ref 93411); α,α' -Azoisobutyronitrile "AIBN" (Fluka, ref 11630); Sodium sulfate (Prolabo; ref 28111.365)

Solvents: Acetonitrile (Riedel-de Haen; ref 33019); Ethyl acetate Pur (Carlo Erba; ref 528299); Dioxan P.A. (Merck; ref 1.09671.1000); Dichloromethane (Merck; ref 1.06050.6025); Methanol (Carlo Erba; ref 309002);

Reference: Robins, M. J., Wilson, J. S., and Hansske, F., Nucleic Acid Related Compounds. 42. A General Procedure for the Efficient Deoxygenation of Secondary Alcohols. Regiospecific and Stereoselective Conversion of Ribonucleosides to 2'-Deoxynucleosides. *J. Am. Chem. Soc.*, **105**, 4059-4065 (1983).

To compound **146** (34 g, 67 mmol) were added acetonitrile (280 ml), DMAP (16.5 g, 135 mmol) and phenyl chlorothionocarbonate (10.2 ml, 73 mmol). The solution was stirred at room temperature for 12 hrs. Solvent was evaporated and the residue was partitioned between ethyl acetate (400 ml) and a HCl 0.5N solution (400 ml). The organic layer was washed with a HCl 0.5N solution (400 ml) and water (2x400 ml), dried over sodium sulfate, filtered and evaporated to dryness to give the intermediate as a pale yellow solid. The crude **147** was dissolved in dioxan (ml) and AIBN (3.3 g, 20 mmol) and TTMSS (33 ml, 107 mmol) were added. The solution was progressively heated until reflux and stirred for 2 hrs. The reaction was concentrated to a yellow oil which was chromatographed (eluent dichloromethane/methanol

95/5) to give compound **148** (23 g, colorless foam, 70%). An aliquot was crystallized from ethanol/ petroleum ether.


Analyses for 3',5'-O-(1,1,3,3-tetraisopropyl-1,3-disiloxanyl)-2'-deoxy- β -L-adenosine **148:**

mp 110-111°C (EtOH/petroleum ether) (Lit.(5) mp 113-114°C (EtOH))

^1H NMR (200 MHz, CDCl_3): δ 8.33 and 8.03 (2s, 2H, H_2 and H_8), 6.30 (dd, 1H, H_1 , J 2.85 Hz, J 7.06 Hz), 5.63 (sl, 2H, NH_2), 4.96 (m, 1H, H_3'), 4.50 (m, 2H, $\text{H}_{5'a}$ and $\text{H}_{5'b}$), 2.68 (m, 2H, $\text{H}_{2'a}$ and $\text{H}_{2'b}$), 1.08 (m, 28H, isopropyl protons)

Mass analysis (FAB+, GT) m/z 494 ($\text{M}+\text{H}$)⁺, 136 (BH_2)⁺

Reaction 5:



Reactants : Ammonium fluoride (Fluka; ref 09742); Silica gel (Merck; ref 1.07734.2500)

Solvents : Methanol P.A. (Prolabo; ref 20847.295); Dichloromethane P.A. (Merck; ref 1.06050.6025); Ethanol 95 (Prolabo; ref 20823.293)

Reference : Zhang, W., and Robins, M. J., Removal of Silyl Protecting Groups from Hydroxyl Functions with Ammonium Fluoride in Methanol. *Tetrahedron Lett.*, **33**, 1177-1180 (192).

A solution of 3',5'-O-(1,1,3,3-tetraisopropyl-1,3-disiloxanyl)-2'-deoxy-L-adenosine **148** (32 g, 65 mmol) and ammonium fluoride (32 g, mmol) in methanol was stirred at reflux for 2 hrs. Silica gel was added and the mixture was carefully evaporated to give a white powder. This powder was added on the top of a silica column, which was eluted with

dichloromethane/methanol 9/1. The appropriate fractions were combined and evaporated to give a white powder, which was crystallized from ethanol 95 (12.1 g, 75%).

Analyses for 2'-Deoxy-β-L-adenosine 149 :

mp 189-190°C (EtOH 95) (identical to commercial 2'-deoxy-D-adenosine)


¹H NMR (200 MHz, DMSO-D₆): δ 8.35 and 8.14 (2s, 2H, H₂ and H₈), 7.34 (sl, 2H, NH₂), 6.35 (dd, 1H, H_{1'}, J 6.1 Hz, J 7.85 Hz), 5.33 (d, 1H, OH_{2'}, J 4.0 Hz), 5.28 (dd, 1H, H_{3'}, J 4.95 Hz; J 6.6 Hz), 4.42 (m, 1H, OH_{5'}), 3.88 (m, 1H, H_{4'}), 3.63-3.52 (m, 2H, H_{5'a} and H_{5'b}), 2.71 (m, 1H, H_{2'a}), 2.28 (m, 1H, H_{2'b}). (identical to commercial 2'-deoxy-D-adenosine)

α_D +26° (c 0.5 water) (commercial 2'-deoxy-D-adenosine -25° (c 0.5 water)).

UV λ_{max} 260 nm (ε 14100) (H₂O).

Mass analysis (FAB+, GT) m/z 252 (M+H)⁺, 136 (BH₂)⁺

Example 3 Stereospecific Synthesis of 2'-Deoxy-β-L-Cytidine



1-(3,5-Di-O-benzoyl-β-L-xylofuranosyl)uracil (**11**)

Hydrazine hydrate (1.4 mL, 28.7 mmol) was added to a solution of 1-(2-O-acetyl-3,5-di-O-benzoyl-β-L-xylofuranosyl)uracil **10** [Ref.: Gosselin, G.; Bergogne, M.-C.; Imbach, J.-L., "Synthesis and Antiviral Evaluation of β-L-Xylofuranosyl Nucleosides of the Five Naturally Occurring Nucleic Acid Bases", Journal of Heterocyclic Chemistry, 1993, 30 (Oct.-Nov.), 1229-1233] (4.79 g, 9.68 mmol) in pyridine (60 mL) and acetic acid (15 mL). The solution was stirred overnight at room temperature. Acetone was added (35 mL) and the mixture was stirred for 30 min. The reaction mixture was evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography [eluent: stepwise gradient of methanol (0-4%) in dichloromethane to give **11** (3.0 g, 68%) which was crystallized from cyclohexane/dichloromethane: mp = 111-114°C; ¹H-NMR (DMSO-*d*₆): δ 11.35 (br s, 1H, NH), 7.9-7.4 (m, 11H, 2 C₆H₅CO, H-6), 6.38 (d, 1H, OH-2', J_{OH-2'} = 4.2 Hz), 5.77 (d, 1H, H-1', J_{1'-2'} = 1.9 Hz), 5.55 (d, 1H, H-5, J₅₋₆ = 8 Hz), 5.54 (dd, 1H, H-3', J_{3'-2'} = 3.9 Hz and J_{3'-4'} = 1.8 Hz), 4.8 (m, 1H, H-4'), 4.7 (m, 2H, H-5' and H-5''), 4.3 (m, 1H, H-2'); MS: FAB>0 (matrix GT) m/z 453 (M+H)⁺, 105 (C₆H₅CO)⁺; FAB<0 (matrix GT) m/z 451 (M-H)⁻, 121 (C₆H₅CO₂)⁻, 111 (B)⁻; Anal. Calcd for C₂₃H₂₀N₂O₈•H₂O : C, 58.09 ; H, 4.76 ; N, 5.96. Found : C, 57.71 ; H, 4.42 ; N, 5.70.

1-(3,5-Di-O-benzoyl-β-L-arabinofuranosyl)uracil (**12**)

To a solution of 1-(3,5-di-O-benzoyl-β-L-xylofuranosyl)uracil **11** (8 g, 17.7 mL) in an anhydrous benzene-DMSO mixture (265 mL, 6:4, v/v) were added anhydrous pyridine (1.4 mL), dicyclohexylcarbodiimide (10.9 g, 53 mmol) and dichloroacetic acid (0.75 mL). The resulting mixture was stirred at room temperature for 4h, then diluted with ethyl acetate (400 mL) and a solution of oxalic acid (4.8 g, 53 mmol) in methanol (14 mL) was added. After being stirred for 1h, the solution was filtered. The filtrate was washed with a saturated NaCl solution (2x500mL), 3% NaHCO₃ solution (2x500mL) and water (2x500mL). The organic phase was dried over Na₂SO₄, then evaporated under reduced pressure. The resulting residue was then solubilized in an EtOH absolute-benzene mixture (140 mL, 2 :1, v/v). To this solution at 0°C was added NaBH₄ (0.96 g, 26.5 mmol). After being stirred for 1h, the solution was diluted with ethyl acetate (400 mL), then filtered. The filtrate was washed with a saturated NaCl solution (400 mL) and water (400 mL). The organic phase was dried over Na₂SO₄, then evaporated under reduced pressure. The resulting crude material was purified by silica gel column chromatography [eluent: stepwise

gradient of methanol (0-3%) in dichloromethane to give **12** (5.3 g, 66%) which was crystallized from acetonitrile: mp = 182-183°C; ¹H-NMR (DMSO-*d*₆): δ 11.35 (br s, 1H, NH), 8.0-7.5 (m, 11H, 2 C₆H₅CO, H-6), 6.23 (br s, 1H, OH-2'), 6.15 (d, 1H, H-1', J_{1'-2'} = 4 Hz), 5.54 (d, 1H, H-5, J₅₋₆ = 8.1 Hz), 5.37 (t, 1H, H-3', J_{3'-2'} = J_{3'-4'} = 2.6 Hz), 4.7-4.6 (m, 2H, H-5' and H-5''), 4.5 (m, 1H, H-4'), 4.4 (m, 1H, H-2'); MS: FAB>0 (matrix GT) m/z 453 (M+H)⁺, 341 (S)⁺, 113 (BH₂)⁺, 105 (C₆H₅CO)⁺; FAB<0 (matrix GT) m/z 451 (M-H)⁻, 121 (C₆H₅CO₂)⁻, 111 (B)⁻; Anal. Calcd for C₂₃H₂₀N₂O₈: C, 61.06 ; H, 4.46 ; N, 6.19. Found : C, 60.83 ; H, 4.34 ; N, 6.25.

1-(3,5-Di-O-benzoyl-2-deoxy-β-L-erythro-pentofuranosyl)uracil (**13**)


To a solution of 1-(3,5-di-O-benzoyl-β-L-arabinofuranosyl)uracil **12** (5.2 g, 11.4 mmol) in anhydrous 1,2-dichloroethane (120 mL) were added phenoxythiocarbonyl chloride (4.7 mL, 34.3 mL) and 4-(dimethylamino)pyridine (DMAP, 12.5 g, 102.6 mmol). The resulting solution was stirred at room temperature under argon atmosphere for 1h and then evaporated under reduced pressure. The residue was dissolved in dichloromethane (300 mL) and the organic solution was successively washed with an ice-cold 0.2 N hydrochloric acid solution (3x 200 mL) and water (2x200 mL), dried over Na₂SO₄ then evaporated under reduced pressure. The crude material was co-evaporated several times with anhydrous dioxane and dissolved in this solvent (110 mL). To the resulting solution were added under argon tris-(trimethylsilyl)silane hydride (4.2 mL, 13.7 mmol) and α,α'-azoisobutyronitrile (AIBN, 0.6 g, 3.76 mmol). The reaction mixture was heated and stirred at 100°C for 1h under argon, then cooled to room temperature and evaporated under reduced pressure. The residue was purified by silica gel column chromatography [eluent: stepwise gradient of methanol (0-5%)] to give **13** (2.78 g, 56%) which was crystallized from EtOH: mp = 223-225°C; H-NMR (DMSO-*d*₆): δ 11.4 (br s, 1H, NH), 8.0-7.5 (m, 11H, 2 C₆H₅CO, H-6), 6.28 (t, 1H, H-1', J = 7 Hz), 5.5 (m, 2H, H-1' and H-5), 4.6-4.4 (m, 3H, H-4', H-5' and H-5''), 2.6 (m, 2H, H-2' and H-2''); MS: FAB>0 (matrix GT) m/z 437 (M+H)⁺, 3325 (S)⁺; FAB<0 (matrix GT) m/z 435 (M-H)⁻, 111 (B)⁻; Anal. Calcd for C₂₃H₂₀N₂O₇: C, 63.30 ; H, 4.62 ; N, 6.42. Found : C, 62.98 ; H, 4.79 ; N, 6.40.

2'-Deoxy-β-L-cytidine (β-L-dC)

Lawesson's reagent (1.72 g, 4.26 mmol) was added under argon to a solution of 1-(3,5-di-O-benzoyl-2-deoxy-β-L-erythro-pentofuranosyl)uracil **13** (2.66 g, 6.1 mmol) in anhydrous 1,2-dichloroethane (120mL) and the reaction mixture was stirred under reflux for 2h. The solvent

was then evaporated under reduced pressure and the residue was purified by silica gel column chromatography [eluent: stepwise gradient of ethyl acetate (0-8%) in dichloromethane] to give the 4-thio intermediate as a yellow foam. A solution of this thio-intermediate (1.5 g, 3.31 mmol) in methanolic ammonia (previously saturated at -10°C and tightly stopped) (50 mL) was heated at 100°C in a stainless-steel bomb for 3h and then cooled to 0°C. The solution was evaporated under reduced pressure. The resulting crude material was purified by silica gel column chromatography [eluent: stepwise gradient of methanol(0-20%) in dichloromethane]. Finally, the appropriate fractions were pooled, filtered through a unit Millex HV-4 (0.45 µm, Millipore) and evaporated under reduced pressure to provide the desired 2'-deoxy-β-L-cytidine (**β-L-dC**) as a foam (0.6 g, 80%) which was crystallized from absolute EtOH: mp=198-199°C; ¹H-NMR (DMSO-*d*₆): δ 7.77 (d, 1H, H-6, *J*₆₋₅ = 7.4 Hz), 7.10 (br d, 2H, NH-2), 6.13 (t, 1H, H-1', *J* = 6.7 Hz), 5.69 (d, 1H, H-5, *J*₅₋₆ = 7.4 Hz), 5.19 (d, 1H, OH-3', *J*_{OH-3'} = 4.1 Hz), 4.96 (t, 1H, OH-5', *J*_{OH-5'} = *J*_{OH-5''} = 5.2 Hz), 4.1 (m, 1H, H-3'), 3.75 (m, 1H, H-4'), 3.5 (m, 2H, H-5' and H-5''), 2.0 (m, 1H, H-2'), 1.9 (m, 1H, H-2''); MS: FAB>0 (matrix GT) *m/z* 228 (M+H)⁺, 112 (BH₂)⁺; FAB<0 (matrix GT) *m/z* 226(M-H)⁻; [α]²⁰_D = - 69 (c 0.52, DMSO) [[α]²⁰_D = + 76 (c 0.55, DMSO) for a commercially available hydrochloride salt of the D-enantiomer]. Anal. Calcd for C₉H₁₃N₃O₄: C, 47.57 ; H, 5.77 ; N, 18.49. Found : C, 47.35 ; H, 5.68 ; N, 18.29.

Example 4 Stereoselective Synthesis of 2'-Deoxy-β-L-Cytidine (β-L-dC)



2-Amino-β-L-arabinofurano[1',2':4,5]oxazoline (**1**)

A mixture of L-arabinose (170 g, 1.13 mol), cyanamide (100g, 2.38 mol), methanol (300 ml), and 6M-NH₄OH (50 ml) was stirred at room temperature for 3 days and then kept at -10°C overnight. The product was collected with suction, washed successively with methanol and ether, and dried *in vacuo*. Yield, 130 g (66.0%) of the analytically pure compound **1**, m.p. 170-172°C; ¹H NMR (DMSO-*d*₆) δ ppm 6.35 (br s, 2H, NH₂), 5.15 (d, 1H, H-1, J= 5.6 Hz), 5.45 (br s, 1H,

OH-3), 4.70 (br s, 1H, OH-5), 4.55 (d, 1H, *H*-2, *J* = 5.6 Hz), 4.00 (br s, 1H, *H*-3), 3.65 (m, 1H, *H*-4), 3.25 (m, 2H, *H*-5, *H*-5').

Reagents:

L-arabinose: Fluka, >99.5%, ref 10839

Cyanamide: Fluka, >98%, ref 28330

***O*^{2,2'}-anhydro-β-L-uridine (**2**)**

A solution of compound **1** (98.8 g, 0.57 mol) and methyl propiolate (98 ml) in 50% aqueous ethanol (740 ml) was refluxed for 5h, then cooled and concentrated under diminished pressure to the half of the original volume. After precipitation with acetone (600 ml), the product was collected with suction, washed with ethanol and ether, and dried. The mother liquor was partially concentrated, the concentrate precipitated with acetone (1000 ml), the solid collected with suction, and washed with acetone and ether to afford another crop of the product. Over-all yield, 80 g (62%) of compound **2**, m.p. 236-240°C; ¹H NMR (DMSO-*d*₆) δ ppm 7.87 (d, 1H, *H*-6, *J* = 7.4 Hz), 6.35 (d, 1H, *H*-1', *J* = 5.7 Hz), 5.95 (d, 1H, *H*-5, *J* = 7.4 Hz), 5.90 (d, 1H, OH-3'), 5.20 (d, 1H, *H*-2', *J* = 5.7 Hz), 5.00 (m, 1H, OH-3'), 4.44 (br s, 1H, *H*-3'), 4.05 (m, 1H, *H*-4'), 3.25 (m, 2H, *H*-5, *H*-5').

Reagent:

Methyl propiolate: Fluka, >97%, ref 81863

3',5'-Di-*O*-benzoyl-*O*^{2,2'}-anhydro-β-L-uridine (3**)**

To a solution of compound **2** (71.1 g, 0.31 mol) in anhydrous pyridine (1200 ml) was added benzoyl chloride (80.4 ml) at 0°C and under argon. The reaction mixture was stirred at room temperature for 5 h under exclusion of atmospheric moisture and stopped by addition of ethanol. The solvents were evaporated under reduced pressure and the resulting residue was coevaporated with toluene and absolute ethanol. The crude mixture was then diluted with ethanol and the precipitate collected with suction, washed successively with ethanol and ether, and dried. Yield, 129 g (95.8%) of compound **3**, m.p. 254 °C; ¹H NMR (DMSO-*d*₆) δ ppm 8.1-7.4 (m, 11H, C₆H₅CO, *H*-6), 6.50 (d, 1H, *H*-1', *J* = 5.7 Hz), 5.90 (d, 1H, *H*-5, *J* = 7.5 Hz), 5.80 (d, 1H, *H*-2', *J*

= 5.8 Hz), 5.70 (d, 1H, H-3') 4.90 (m, 1H, H-4'), 4.35 (m, 2H, H-5, H-5').

Reagent:

Benzoyl chloride: Fluka, p.a., ref 12930

3',5'-Di-O-benzoyl-2'-chloro-2'-deoxy- β ,L-uridine (4)

To a solution of compound 3 (60.3 g, 0.139 mol) in dimethylformamide (460 ml) was added at 0°C a 3.2 N-HCl/DMF solution (208 ml, prepared *in situ* by adding 47.2 ml of acetyl chloride at 0°C to a solution of 27.3 ml of methanol and 133.5 ml of dimethylformamide). The reaction mixture was stirred at 100°C for 1h under exclusion of atmospheric moisture, cooled down, and poured into water (4000 ml). The precipitate of compound 4 was collected with suction, washed with water, and recrystallised from ethanol. The crystals were collected, washed with cold ethanol and ether, and dried under diminished pressure. Yield, 60.6 g (92.6%) of compound 4, m.p. 164-165°C; ¹H NMR (DMSO-*d*₆) δ ppm 8.7 (br s, 1H, NH), 8.1-7.3 (m, 11H, C₆H₅CO, H-6), 6.15 (d, 1H, H-1', J = 4.8 Hz), 5.5 (m, 2H, H-5, H-2'), 4.65 (m, 4H, H-3', H-4', H-5', H-5'').

Reagent:

Acetyl chloride: Fluka, p.a., ref 00990

3',5'-Di-O-benzoyl-2'-deoxy- β ,L-uridine (5)

A mixture of compound 4 (60.28 g, 0.128 mol), tri-n-butyltin hydride (95 ml) and azabisisobutyronitrile (0.568 g) in dry toluene (720 ml) was refluxed under stirring for 5 h and cooled down. The solid was collected with suction and washed with cold toluene and petroleum ether. The filtrate was concentrated under reduced pressure and diluted with petroleum ether to deposit an additional crop of compound 5. Yield, 54.28 g (97.2%) of compound 5; m.p. 220-221°C ; ¹H NMR (CDCl₃) δ ppm 8.91 (br s, 1H, NH), 8.1-7.5 (m, 11H, C₆H₅CO and H-6), 6.43 (q, 1H, H-1', J_{1',2'} = 5.7 Hz and J_{1',2''} = 8.3 Hz), 5.7-5.6 (m, 2H, H-3' and H-5), 4.8-4.6 (m, 3H, H-5', H-5'' and H-4'), 2.8-2.7 (m, 1H, H-2'), 2.4-2.3 (m, 1H, H-2'').

Reagents:

Tri-n-butyltin hydride: Fluka, >98%, ref 90915

Azabisisobutyronitrile: Fluka, >98%, ref 11630

3',5'-Di-*O*-benzoyl-2'-deoxy- β -L-4-thio-uridine (6)

A solution of compound 5 (69 g, 0.158 mol) and Lawesson's reagent (74 g) in anhydrous methylene chloride (3900 ml) was refluxed under argon overnight. After evaporation of the solvent, the crude residue was purified by a silica gel column chromatography [eluant: gradient of methanol (0-2%) in methylene chloride] to afford pure compound 6 (73 g) in quantitative yield; ^1H NMR (CDCl_3) δ ppm 9.5 (br s, 1H, *NH*), 8.1-7.4 (m, 10H, $\text{C}_6\text{H}_5\text{CO}$), 7.32 (d, 1H, *H*-6, $J = 7.7$ Hz), 6.30 (dd, 1H, *H*-1', $J = 5.6$ Hz and $J = 8.2$ Hz), 6.22 (d, 1H, *H*-5, $J = 7.7$ Hz), 5.6 (m, 1H, *H*-3'), 4.7 (m, 2H, *H*-5', *H*-5''), 4.5 (m, 1H, *H*-4'), 2.8 (m, 1H, *H*-2'), 2.3 (m, 1H, *H*-2'').

Reagent:

Lawesson's reagent: Fluka, >98%, ref 61750


2'-Deoxy- β -L-cytosine

A solution of compound 6 (7.3 g, 0.016 mol) in methanol saturated with ammonia (73 ml) was heated at 100°C in a stainless steel cylinder for 3h. After cooling carefully, the solvent was evaporated under reduced pressure. An aqueous solution of the residue was washed with ethyl acetate and evaporated to dryness. Such a procedure was carried out on 9 other samples (each 7.3 g) of compound 6 (total amount of 6 = 73 g). The 10 residues were combined, diluted with absolute ethanol and cooled to give 7 as crystals. Trace of benzamide were eliminated from the crystals of 6 by a solid-liquid extraction procedure (at reflux in ethyl acetate for 1h). Yield, 28.75 g (78.6%) of compound 6; m. p. 141-145°C ; ^1H NMR (DMSO) δ ppm 8.22 and 8.00 (2 br s, 2H, NH_2), 7.98 (d, 1H, *H*-6, $J = 7.59$ Hz), 6.12 (t, 1H, *H*-1', $J = 6.5$ Hz and $J = 7.6$ Hz), 5.89 (d, 1H, *H*-5, $J = 7.59$ Hz), 5.3 (br s, 1H, *OH*-3'), 5.1 (br s, 1H, *OH*-5'), 4.2 (m, 1H, *H*-3'), 3.80 (q, 1H, *H*-4', $J = 3.6$ Hz and $J = 6.9$ Hz), 3.6-3.5 (m, 2H, *H*-5', *H*-5''), 2.2-2.0 (m, 2H, *H*-2', *H*-2''); FAB<0, (GT) m/e 226 (M-H)⁻, 110 (B)⁻; FAB>0 (GT) 228 (M+H)⁺, 112 (B+2H)⁺; $[\alpha]_{\text{D}}^{20}$ - 56.48 ($c = 1.08$ in DMSO); UV (pH 7) $\lambda_{\text{max}} = 270$ nm ($\epsilon = 10000$).

Reagent:

Methanolic ammonia: previously saturated at -5°C, tightly stoppered, and kept in a freezer.

Example 5 Stereoselective Synthesis of 2'-Deoxy-β-L-Thymidine (β-L-dT)



3',5'-Di-O-benzoyl-2'-deoxy-5-iodo-β-L-uridine (7)

A mixture of compound **5** (105.8 g, 0.242 mol), iodine (76.8 g), CAN (66.4 g) and acetonitrile (2550 ml) was stirred at 80°C for 3h then the reaction mixture was cooled at room temperature leading to crystallization of compound **7** (86.6 g, 63.5%); m. p. 192-194°C ; ^1H NMR (DMSO) δ ppm .8.34 (s, 1H, NH), 8.2-7.2 (m, 11H, 2 C₆H₅CO, H-6), 6.31 (q, 1H, H-1', J = 5.5 Hz and J = 8.7 Hz), 5.5 (m, 1H, H-3'), 4.7 (m, 2H, H-5', H-5''), 4.5 (m, 1H, H-4'), 2.7 (m, 1H, H-2'), 2.3 (m, 1H, H-2''); FAB<0, (GT) m/e 561 (M-H)⁻, 237 (B)⁻; FAB>0 (GT) 563 (M+H)⁺; $[\alpha]_{\text{D}}^{20} + 39.05$ (c = 1.05 in DMSO); UV (EtOH 95) ν_{max} = 281 nm (ϵ = 9000), ν_{min} = 254 nm (ϵ = 4000), ν_{max} = 229 nm (ϵ = 31000); Anal. Calcd for C₂₃H₁₉IN₂O₇: C, 49.13 H, 3.41 N, 4.98 I, 22.57. Found: C, 49.31 H, 3.53 N, 5.05 I, 22.36.

Reagents:

Iodine: Fluka, 99.8%, ref 57650

Cerium ammonium nitrate (CAN): Aldrich, >98.5%, ref 21,547-3

3',5'-Di-*O*-benzoyl-2'-deoxy-3-*N*-toluoyl- β -L-thymidine (9**)**

To a solution of compound **7** (86.6g, 0.154 mol) in anhydrous pyridine (1530 ml) containing *N*-ethyldiisopropylamine (53.6 ml) was added, portionwise at 0°C, *p*-toluoyl chloride (40.6 ml). The reaction mixture was stirred for 2 h at room temperature, then water was added to stop the reaction and the reaction mixture was extracted with methylene chloride. The organic phase was washed with water, dried over sodium sulfate and evaporated to dryness to give crude 3',5'-di-*O*-benzoyl-2'-deoxy-3-*N*-toluoyl-5-iodo- β -L-uridine (**8**) which can be used for the next step without further purification.

A solution of the crude mixture **8**, palladium acetate (3.44 g), triphenylphosphine (8.0 g) in *N*-methylpyrrolidinone (1375 ml) with triethylamine (4.3 ml) was stirred at room temperature for 45 min. Then, tetramethyltin (42.4 ml) was added dropwise at 0°C under argon. After stirring at 100-110°C overnight, the reaction mixture was poured into water and extracted with diethyl ether. The organic solution was dried over sodium sulfate and concentrated under reduced pressure. The residue was purified by a silica gel column chromatography [eluant: stepwise gradient of ethyl acetate (0-10%) in toluene] to give compound **9** as a foam (42.3 g, 48.3% for the 2 steps). ¹H NMR (DMSO) δ ppm 8.3-7.2 (m, 15H, 2 C₆H₅CO, 1 CH₃C₆H₄CO, H-6), 6.29 (t, 1H, *H*-1', *J* = 7.0 Hz), 5.7 (m, 1H, *H*-3'), 4.7-4.5 (m, 3H, *H*-5', *H*-5'', *H*-4'), 2.7-2.6 (m, 2H, *H*-2', *H*-2''); FAB<0, (GT) *m/e* 567 (M-H)⁻, 449 (M-CH₃C₆H₄CO)⁻, 243 (B)⁻, 121 (C₆H₅COO)⁻; FAB>0 (GT) 1137 (2M+H)⁺, 569 (M+H)⁺, 325 (M-B)⁻, 245 (B+2H)⁺, 119 (CH₃C₆H₅CO)⁺.

Reagents:

***p*-Toluoyl chloride,** Aldrich, 98%, ref 10,663-1

Diisopropylethylamine: Aldrich, >99.5%, ref 38,764-9

***N*-methylpyrrolidinone:** Aldrich, >99%, ref 44,377-8

Palladium acetate: Aldrich, >99.98%, ref 37,987-5

Triphenylphosphine: Fluka, >97%, ref 93092

Tetramethyltin: Aldrich, >99%, ref 14,647-1

2'-Deoxy- β -L-thymidine


A solution of compound **9** (42.3 g, 0.074 mol) in methanol saturated with ammonia (1850 ml) was stirred at room temperature for two days. After evaporation of the solvent, the residue was diluted with water and washed several times with ethyl acetate. The aqueous layer was separated, evaporated under reduced pressure and the residue was purified by a silica gel column chromatography [eluant: stepwise gradient of methanol (0-10%) in methylene chloride] to give pure 2'-deoxy- β -L-thymidine (11.62 g, 64.8%) which was crystallized from ethanol; m.p. 185-188°C; ^1H NMR (DMSO) δ ppm 11.3 (s, 1H, NH), 7.70 (s, 1H, *H*-6), 6.2 (pt, 1H, *H*-1'), 5.24 (d, 1H, OH-3', *J* = 4.2 Hz), 5.08 (t, 1H, OH-5', *J* = 5.1 Hz), 4.2 (m, 1H, *H*-3'), 3.7 (m, 1H, *H*-4'), 3.5-3.6 (m, 2H, *H*-5', *H*-5''), 2.1-2.0 (m, 2H, *H*-2', *H*-2''); FAB<0, (GT) *m/e* 483 (2M-H)⁻, 349 (M+T-H)⁻, 241 (M-H)⁻, 125 (B)⁻; FAB>0 (GT) 243 (M+H)⁺, 127 (B+2H)⁺; $[\alpha]_{\text{D}}^{20}$ - 13.0 (*c* = 1.0 in DMSO); UV (pH 1) ν_{max} = 267 nm (ϵ = 9700), ν_{min} = 234 nm (ϵ = 2000).

Reagent:

Methanolic ammonia: previously saturated at -5°C, tightly stoppered, and kept in a freezer.

Example 6 Stereoselective Synthesis of 2'-deoxy- β -L-inosine (β -L-dI)

β -L-dI was synthesized by deamination of 2'-deoxy- β -L-adenosine (β -L-dA) following a procedure previously described in the 9-D-glucopyranosyl series (Ref: I. Iwai, T. Nishimura and B. Shimizu, Synthetic Procedures in Nucleic Acid Chemistry, W. W. Aorbach and R. S. Tipson, eds., John Wiley & Sons, Inc. New York, vol. 1, pp. 135-138 (1968)).



Thus, a solution of $\beta\text{-L-dA}$ (200 mg) in a mixture of acetic acid (0.61 ml) and water (19 ml) was heated with sodium nitrite (495 mg), and the mixture was stirred at room temperature overnight. The solution was then evaporated to dryness under diminished pressure. An aqueous solution of the residue was applied to a column of IR-120 (H^+) ion-exchange resin, and the column was eluted with water. Appropriate fractions were collected and evaporated to dryness to afford pure $\beta\text{-L-dI}$ which was crystallized from methanol (106 mg, 53% yield not optimized): m.p.=209°-211°C; UV (H_2O), λ_{max} =247 nm; $^1\text{H-NMR}$ (DMSO-d_6)= 8.32 and 8.07 (2s, 1H each, H-2 and H-8), 6.32 (pt, 1H, H-1; $J=6.7$ Hz), 4.4 (m, 1H, H-3'), 3.9 (m, 1H, H-4'), 3.7-3.4 (m, 2H partially obscured by HOD, H-5',5''), 2.6 and 2.3 (2m, 1H each, H-2' and H-2''); mass spectra (mature, glycerol-thioglycerol, 1:1, v/v), FAB>0: 253 ($m+\text{H}$)⁺, 137 (base + 2H)⁺; FAB<0: 251 ($m-\text{H}$)⁻, 135 (base)⁻; $[\alpha]_{\text{D}}^{20} = +19.3$ (-c 0.88, H_2O).

Anti-HBV Activity of the Active Compounds

The ability of the active compounds to inhibit the growth of virus in 2.2.15 cell cultures (HepG2 cells transformed with hepatitis virion) can be evaluated as described in detail below.

A summary and description of the assay for antiviral effects in this culture system and the analysis of HBV DNA has been described (Korba and Milman, 1991, *Antiviral Res.*, 15:217). The antiviral evaluations are performed on two separate passages of cells. All wells, in all plates, are seeded at the same density and at the same time.

Due to the inherent variations in the levels of both intracellular and extracellular HBV DNA, only depressions greater than 3.5-fold (for HBV virion DNA) or 3.0-fold (for HBV DNA replication intermediates) from the average levels for these HBV DNA forms in untreated cells are considered to be statistically significant ($P<0.05$). The levels of integrated HBV DNA in each

cellular DNA preparation (which remain constant on a per cell basis in these experiments) are used to calculate the levels of intracellular HBV DNA forms, thereby ensuring that equal amounts of cellular DNA are compared between separate samples.

Typical values for extracellular HBV virion DNA in untreated cells range from 50 to 150 pg/ml culture medium (average of approximately 76 pg/ml). Intracellular HBV DNA replication intermediates in untreated cells range from 50 to 100 $\mu\text{g/pg}$ cell DNA (average approximately 74 pg/ μg cell DNA). In general, depressions in the levels of intracellular HBV DNA due to treatment with antiviral compounds are less pronounced, and occur more slowly, than depressions in the levels of HBV virion DNA (Korba and Milman, 1991, Antiviral Res., 15:217).

The manner in which the hybridization analyses are performed for these experiments result in an equivalence of approximately 1.0 pg of intracellular HBV DNA to 2-3 genomic copies per cell and 1.0 pg/ml of extracellular HBV DNA to 3×10^5 viral particles/ml.

Example 7

The ability of the triphosphate derivatives of $\beta\text{-L-dA}$, $\beta\text{-L-dC}$, $\beta\text{-L-dU}$, $\beta\text{-L-2'-dG}$, $\beta\text{-L-dI}$, and $\beta\text{-L-dT}$ to inhibit hepatitis B was tested. Table 1 describes the comparative inhibitory activities of triphosphates of $\beta\text{-L-dT}$ ($\beta\text{-L-dT-TP}$), $\beta\text{-L-dC}$ ($\beta\text{-L-dC-TP}$), $\beta\text{-L-dU}$ ($\beta\text{-L-dU-TP}$) and $\beta\text{-L-dA}$ ($\beta\text{-L-dA-TP}$) on woodchuck hepatitis virus (WHV) DNA polymerase, human DNA polymerases α , β , and γ .

Table 1

Inhibitor	WHV DNA pol IC ₅₀	DNA pol α K _i ^b (μ M)	DNA pol β K _i ^b (μ M)	DNA pol γ K _i ^b (μ M)
β -L-dT-TP	0.34	>100	>100	>100
β -L-dA-TP	2.3	>100	>100	>100
β -L-dC-TP	2.0	>100	>100	>100
β -L-dU-TP	8	>100	>100	>100

^aIC₅₀: 50% Inhibitory concentration

^bK_i value was determined using calf thymus activated DNA as template-primer and dATP as substrate. Inhibitors were analyzed by Dixon plot analysis. Under these conditions, the calculated mean K_m of human DNA polymerase α for dATP as approximately 2.6 μ M. Human DNA polymerase β exhibited a steady state K_m of 3.33 μ M for dATP. Human DNA polymerase γ exhibited a steady K_m of 5.2 μ M.

Example 8

The anti-hepatitis B virus activity of β -L-dA, β -L-dC, β -L-dU, β -L-2'-dG and β -L-dT was tested in transfected Hep G-2 (2.2.15) cells. Table 2 illustrates the effect of β -L-dA, β -L-dC, β -L-dU, and β -L-dT against hepatitis B virus replication in transfected Hep G-2 (2.2.15) cells.

Table 2

Compound	HBV virions ^a EC ₅₀ (μ M)	HBV Ri ^b EC ₅₀ (μ M)	Cytotoxicity IC ₅₀ (μ M)	Selectivity Index IC ₅₀ /EC ₅₀
β -L-dT	0.05	0.05	>200	>4000
β -L-dC	0.05	0.05	>200	>4000
β -L-dA	0.10	0.10	>200	>2000
β -L-dI	1.0	1.0	>200	>200
β -L-dU	5.0	5.0	>200	>40

^aExtracellular DNA

^bReplicative intermediates (Intracellular DNA)

Example 9

The effect of β -L-dA, β -L-dC and β -L-dT in combination on the growth of hepatitis B was measured in 2.2.15 cells. The results are provided in Table 3.

Table 3

Combination	Ratio	EC ₅₀
L-dC + L-dT	1:3	.023
L-dC + L-dT	1:1	.053
L-dC + L-dT	3:1	.039
L-dC + L-dA	1:30	.022
L-dC + L-dA	1:10	.041
L-dC + L-dA	1:3	.075
L-dT + L-dA	1:30	.054
L-dT + L-dA	1:10	.077
L-dT + L-dA	1:3	.035

Each combination produced anti-HBV activity that was synergistic. In addition, the combination of L-dA + L-dC + L-dT was also synergistic in this model.

Example 10

The inhibition of hepatitis B replication in 2.2.15 cells by β -L-dA and β -L-dC, alone and in combination was measured. The results are shown in Table 4.

Table 4

^a β-L-2'-deoxy-adenosine (μM)	^b β-L-2'-deoxy-cytidine (μM)	% Inhibition	^c C.I.
0.5		90	
0.05		24	
0.005		1	
	0.5	95	
	0.05	40	
	0.005	10	
0.05	0.05	80	0.34
0.05	0.005	56	0.20
0.05	0.0005	50	0.56
0.005	0.05	72	0.35
0.005	0.005	54	0.35
0.005	0.0005	30	0.16
0.0005	0.05	50	0.83
0.0005	0.005	15	0.28
0.0005	0.0005	0	N.A.

^aβ-L-2'-deoxy-adenosine: IC₅₀=0.09 μM

^bβ-L-2'-deoxy-cytidine: IC₅₀=0.06 μM

^cCombination indices values indicate synergism effect (<1), additive effect (=1), and antagonism effect (>1)

Example 11

The efficacy of L-dA, L-dT and L-dC against hepadnavirus infection in woodchucks (*Marmota monax*) chronically infected with woodchuck hepatitis virus (WHV) was determined. This animal model of HBV infection is widely accepted and has proven to be useful for the evaluation of antiviral agents directed against HBV.

Protocol:

Experimental groups (n=3 animals/drug group, n=4 animals/control)

Group 1	vehicle control
Group 2	lamivudine (3TC) (10 mg/kg/day)
Groups 3-6	L-dA (0.01, 0.1, 1.0, 10 mg/kg/day)
Groups 7-10	L-dT (0.01, 0.1, 1.0, 10 mg/kg/day)
Groups 11-14	L-dC (0.01, 0.1, 1.0, 10 mg/kg/day)

Drugs were administered by oral gavage once daily, and blood samples taken on days 0, 1, 3, 7, 14, 21, 28, and on post-treatment days +1, +3, +7, +14, +28 and +56. Assessment of the activity and toxicity was based on the reduction of WHV DNA in serum: dot-blot, quantitative PCR.

The results are illustrated in Figure 3 and Table 5.

Table 5
Antiviral Activity of LdA, LdT, LdC in Woodchuck Model of Chronic HBV Infection

day	Control	LdA	LdT	LdC
	ng WHV-DNA per ml serum ^{1,2}			
0	381	436	423	426
1	398	369	45	123
3	412	140	14	62
7	446	102	6	46
14	392	74	1	20

¹ LdA, LdT, LdC administered orally once a day at 10mg/kg

² limit of detection is 1 ng/ml WHV-DNA per ml serum

The data show that L-dA, L-dT and L-dC are highly active in this in vivo model. First, viral load is reduced to undetectable (L-dT) or nearly undetectable (L-dA, L-dC) levels. Second, L-dA, L-dT and L-dC are shown to be more active than 3TC (lamivudine) in this model. Third, viral rebound is not detected for at least two weeks after withdrawal of L-dT. Fourth, dose response curves suggest that a modest increase in the dose of L-dA and L-dC would show antiviral activity similar to L-dT. Fifth, all animals receiving the drugs gained weight and no drug-related toxicity was detected.

Toxicity of Compounds

Toxicity analyses were performed to assess whether any observed antiviral effects are due to a general effect on cell viability. The method used is the measurement of the effect of β -L-dA, β -L-dC and β -L-dT on cell growth in human bone marrow clonogenic assays, as compared to Lamivudine. The results are provided in Table 6.

Table 6

Compound	CFU-GM (μ M)	BFU-E (μ M)
β -L-dA	>10	>10
β -L-dC	>10	>10
β -L-dT	>10	>10
β -L-dU	>10	>10
Lamuvudine	>10	>10

Preparation of Pharmaceutical Compositions

Humans suffering from any of the disorders described herein, including hepatitis B, can be treated by administering to the patient an effective amount of a β -2'-deoxy- β -L-erythro-pentofuranonucleoside, for example, β -L-2'-deoxyadenosine, β -L-2'-deoxycytidine, β -L-2'-deoxyuridine, β -L-2'-deoxyguanosine or β -L-2'-deoxythymidine or a pharmaceutically acceptable prodrug or salt thereof in the presence of a pharmaceutically acceptable carrier or diluent. The active materials can be administered by any appropriate route, for example, orally, parenterally, intravenously, intradermally, subcutaneously, or topically, in liquid or solid form.

The active compound is included in the pharmaceutically acceptable carrier or diluent in an amount sufficient to deliver to a patient a therapeutically effective amount of compound to inhibit viral replication *in vivo*, without causing serious toxic effects in the patient treated. By "inhibitory amount" is meant an amount of active ingredient sufficient to exert an inhibitory effect as measured by, for example, an assay such as the ones described herein.

A preferred dose of the compound for all of the abovementioned conditions will be in the range from about 1 to 50 mg/kg, preferably 1 to 20 mg/kg, of body weight per day, more generally 0.1 to about 100 mg per kilogram body weight of the recipient per day. The effective dosage range of the pharmaceutically acceptable prodrug can be calculated based on the weight of the parent nucleoside to be delivered. If the prodrug exhibits activity in itself, the effective

dosage can be estimated as above using the weight of the prodrug, or by other means known to those skilled in the art.

The compound is conveniently administered in unit any suitable dosage form, including but not limited to one containing 7 to 3000 mg, preferably 70 to 1400 mg of active ingredient per unit dosage form. A oral dosage of 50-1000 mg is usually convenient.

Ideally the active ingredient should be administered to achieve peak plasma concentrations of the active compound of from about 0.2 to 70 μM , preferably about 1.0 to 10 μM . This may be achieved, for example, by the intravenous injection of a 0.1 to 5% solution of the active ingredient, optionally in saline, or administered as a bolus of the active ingredient.

The concentration of active compound in the drug composition will depend on absorption, inactivation, and excretion rates of the drug as well as other factors known to those of skill in the art. It is to be noted that dosage values will also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition. The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at varying intervals of time.

A preferred mode of administration of the active compound is oral. Oral compositions will generally include an inert diluent or an edible carrier. They may be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition.

The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent

such as peppermint, methyl salicylate, or orange flavoring. When the dosage unit form is a capsule, it can contain, in addition to material of the above type, a liquid carrier such as a fatty oil. In addition, dosage unit forms can contain various other materials which modify the physical form of the dosage unit, for example, coatings of sugar, shellac, or other enteric agents.

The compound can be administered as a component of an elixir, suspension, syrup, wafer, chewing gum or the like. A syrup may contain, in addition to the active compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors.

The compound or a pharmaceutically acceptable derivative or salts thereof can also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action, such as antibiotics, antifungals, antiinflammatories, protease inhibitors, or other nucleoside or nonnucleoside antiviral agents. Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parental preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

If administered intravenously, preferred carriers are physiological saline or phosphate buffered saline (PBS).

In a preferred embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation.


Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) are also preferred as pharmaceutically acceptable carriers. These

may be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811 (which is incorporated herein by reference in its entirety). For example, liposome formulations may be prepared by dissolving appropriate lipid(s) (such as stearyl phosphatidyl ethanolamine, stearyl phosphatidyl choline, arachadoyl phosphatidyl choline, and cholesterol) in an inorganic solvent that is then evaporated, leaving behind a thin film of dried lipid on the surface of the container. An aqueous solution of the active compound or its monophosphate, diphosphate, and/or triphosphate derivatives is then introduced into the container. The container is then swirled by hand to free lipid material from the sides of the container and to disperse lipid aggregates, thereby forming the liposomal suspension.

This invention has been described with reference to its preferred embodiments. Variations and modifications of the invention, will be obvious to those skilled in the art from the foregoing detailed description of the invention. It is intended that all of these variations and modifications be included within the scope of the this invention.


We claim:

1. A method for treating a host infected with hepatitis B comprising administering an effective amount of an anti-HBV biologically active 2'-deoxy- β -L-erythro-pentofuranonucleoside or a pharmaceutically acceptable salt or prodrug thereof, wherein the 2'-deoxy- β -L-erythro-pentofuranonucleoside has the formula:




wherein R is selected from the group consisting of H, straight chained, branched or cyclic alkyl, CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, amino acid residue, mono, di, or triphosphate, or a phosphate derivative; and BASE is a purine or pyrimidine base which may be optionally substituted.

2. The method of claim 1, wherein the 2'-deoxy- β -L-erythro-pentofuranonucleoside is β -L-2'-deoxyadenosine or a pharmaceutically acceptable salt or prodrug thereof, of the formula:




wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

3. The method of claim 1, wherein the 2'-deoxy- β -L-erythro-pentofuranonucleoside is β -L-2'-deoxycytidine or pharmaceutically acceptable salt or prodrug thereof of the formula:




wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

4. The method of claim 1, wherein the 2'-deoxy- β -L-erythro-pentofuranonucleoside is β -L-2'-deoxyuridine or pharmaceutically acceptable salt or prodrug thereof of the formula:




wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

5. The method of claim 1, wherein the 2'-deoxy- β -L-erythro-pentofuranonucleoside is β -L-2'-deoxyguanosine or pharmaceutically acceptable salt or prodrug thereof of the formula:




wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

6. The method of claim 1, wherein the 2'-deoxy- β -L-erythro-pentofuranonucleoside is β -L-2'-deoxyinosine or pharmaceutically acceptable salt or prodrug thereof of the formula:




wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

7. The method of claim 1, wherein the 2'-deoxy- β -L-erythro-pentofuranonucleoside is β -L-thymidine or a pharmaceutically acceptable salt or prodrug thereof of the formula:




wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

8. A method for treating a host infected with hepatitis B comprising administering an effective amount of two or more anti-HBV biologically active 2'-deoxy- β -L-erythro-pentofuranonucleosides or a pharmaceutically acceptable salt or prodrug thereof in combination or alternation, wherein the 2'-deoxy- β -L-erythro-pentofuranonucleosides have the formula:



wherein R is selected from the group consisting of H, straight chained, branched or cyclic alkyl, CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, amino acid residue, mono, di, or triphosphate, or a phosphate derivative; and BASE is a purine or pyrimidine base which may be optionally substituted.


9. A method for treating a host infected with hepatitis B comprising administering an effective amount of an anti-HBV biologically active 2'-deoxy-β-L-erythro-pentofuranonucleoside or a pharmaceutically acceptable salt or prodrug thereof in combination or alternation with an additional anti-hepatitis B agent, wherein the 2'-deoxy-β-L-erythro-pentofuranonucleoside has the formula:



wherein R is selected from the group consisting of H, straight chained, branched or cyclic alkyl, CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, amino acid residue, mono, di, or triphosphate, or a phosphate derivative; and BASE is a purine or pyrimidine base which may be optionally substituted.

10. The method of claim 9, wherein the additional anti-hepatitis B agent is selected from the group consisting of 3TC, FTC, L-FMAU, DAPD, famciclovir, penciclovir, BMS-200475, bis pom PMEA (adefovir, dipivoxil); lobucavir, ganciclovir, and ribavarin

11. A compound or pharmaceutically acceptable salt or prodrug thereof of the formula:




wherein R is H, mono, di or tri phosphate, acyl, or alkyl, or a stabilized phosphate derivative (to form a stabilized nucleotide prodrug).

12. A pharmaceutical composition comprising an effective amount of a compound of claim 11 in combination with a pharmaceutically acceptable carrier.

Abstract of the Disclosure

This invention is directed to a method for treating a host infected with hepatitis B comprising administering an effective amount of an anti-HBV biologically active 2'-deoxy- β -L-erythro-pentofuranonucleoside or a pharmaceutically acceptable salt or prodrug thereof, wherein the 2'-deoxy- β -L-erythro-pentofuranonucleoside has the formula:



wherein R is selected from the group consisting of H, straight chained, branched or cyclic alkyl, CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, amino acid residue, mono, di, or triphosphate, or a phosphate derivative; and BASE is a purine or pyrimidine base which may be optionally substituted. The 2'-deoxy- β -L-erythro-pentofuranonucleoside or a pharmaceutically acceptable salt or prodrug thereof may be administered either alone or in combination with another 2'-deoxy- β -L-erythro-pentofuranonucleoside or in combination with another anti-hepatitis B agent.

Process for Preparing β -L-d-Nucleosides




Figure 1

Accumulation and Decay of L-dA, L-dT, and L-dC in HepG2 Cells




Figure 2

Antiviral Effect of β L-dA, β L-dT and β L-dC in Woodchuck Chronic Hepatitis Model

(n=3 per drug treatment group, n=4 per placebo group,
dose 10 mg/kg orally once per day)




Figure 3

DECLARATION FOR PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name;

I believe I am the original, first and sole inventor (if only one name is listed below), or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

β -L-2'-Deoxy-Nucleosides for the Treatment of Hepatitis B Virus

the specification of which (check one)

 x is attached hereto
_____ was filed on _____
as application Serial No. _____
_____ and was amended on: _____
(if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability and examination of this application in accordance with Title 37, Code of Federal Regulations, § 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, § 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority
Claimed

(Number)

(Country)

(Day/Month/Year Filed)

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of the application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No.	Filing Date	Status
60/096,110	August 10, 1998	pending
60/131,352	April 28, 1999	pending

As named inventor, I hereby appoint the following attorney(s) or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Sherry M. Knowles, Reg. No. 33,052
Jacqueline Haley, Reg. No. 41,457
Curtis L. Doster, Reg. No. 41,714

Send Correspondence to: Sherry M. Knowles, Esq.
King & Spalding
191 Peachtree Street
Atlanta, Georgia 30303-1763

Direct telephone calls to: Sherry M. Knowles (404) 572-3541

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor Gilles Gosselin

Inventor's signature _____ Date _____

Residence 82 Rue Calvin, 400 Avenue Paul Rimbaud, 34080 Montpellier France

Citizenship France

Post Office Address Same as above

Full name of second joint inventor Jean-Louis Imbach

Inventor's signature _____ Date _____

Residence 1108 Reu las Sorbes, Impasse des Luquesop, 34000 Montpellier, France

Citizenship France

Post Office Address Same as above

Full name of third joint inventor Martin L. Bryant

Inventor's signature _____ Date _____

Residence 65 Hickory Lane, Carlisle, MA 01741

Citizenship U.S.A.

Post Office Address Same as above

Full name of fourth joint inventor _____

Inventor's signature _____ Date _____

Residence _____

Citizenship _____

Post Office Address Same as above